

Notes on duality theories of abelian groups ¹

Gábor Lukács ²

February 2, 2008

¹2000 Mathematics Subject Classification: 22-02 (22A05 43A40)

²I gratefully acknowledge the generous financial support received from the Killam Trusts and Dalhousie University that enabled me to do this research.

Contents

Introduction	1
I The Pontryagin dual	3
I.A The evaluation homomorphism	3
I.1 Pontryagin dual	3
I.2 Polar	3
I.6 Evaluation homomorphism	5
I.10 Local quasi-convexity	7
I.21 Precompactness	10
I.24 Quasi-convex compactness	10
I.B Special groups and subgroups	11
I.28 Metrizable groups	11
I.35 Compact and discrete groups	14
I.38 Bohr-compactification	15
I.41 Open and compact subgroups	16
I.47 Locally compact groups	19
II Categorical and topological generalizations	22
II.A Group objects in a category	22
II.1 Defining groups using diagrams	22
II.2 $\text{Grp}(\mathfrak{X})$ and $\text{Ab}(\mathfrak{X})$	22
II.3 Cartesian closedness	22
II.4 Internal hom	22
II.B k -groups	22
II.5 $\text{Grp}(k\text{Haus})$	22
II.6 k -groups of Noble	22
II.C Convergence groups	23
Appendix	24
A Separation properties of topological groups	24
B Exponentiability and the compact-open topology	25
C Hausdorff k -spaces	27

Introduction

In December 2005, I had the good fortune of spending a week as the guest of Reinhard Winkler at Technische Universität Wien, and had the honour of being asked to deliver a mini-course on the topic of duality theory of abelian groups. During this visit, I came to realize that the only elementary text on this topic that I could refer a graduate student to, namely the one by Pontryagin [30], was published 20 years ago, and its well-organized content is half a century old. This is not to say that there are no excellent books that contain a lot of results on abelian topological groups (such as [12], [33], [26], or [2]), but my feeling is that they depart from the aim of presenting basic notions of duality theory in a self-contained and elementary manner. Therefore, encouraged by my host and his research group in Vienna, I decided to write up the notes that I prepared for the mini course (Chapters 1 and 2), and to keep developing it as a longer term project. In order to provide a smoother presentation, some elementary results of were collected in the Appendix.

I use *italics font* for results that appear to be new and not part of the "common knowledge," while *slanted font* is used for all other statements. Let me know if I am wrong about any of them.

Past and future work

This notebook is under development, so all suggestions, comments, and questions are warmly welcome.

Chapter 1. This is definitely the only more-or-less ready piece, but I am not happy with my treatment of locally compact abelian groups. (Similarly to Roeder's approach (cf. [32]), the only result borrowed from functional analysis is the Peter-Weyl theorem. I am still missing a nice proof of the structure theorem of compactly generated LCA groups, and a complete proof of the Pontryagin duality.)

Chapter 2. I have only a sketch. After a categorical introduction, which explains why cartesian closed categories are so interesting, I hope to cover k -groups (of Noble) and convergence groups in this chapter.

Appendix. It evolves as the chapters develop. It is a collection of results many readers might be familiar with, so I saw no point in including them into the chapters.

Future topics. Precompact (abelian) groups [the problem is that it requires the reader to be familiar with notions such as C -embedded subsets, etc]; examples of pathological or otherwise interesting groups [although [12] is probably the best source for this]; localic groups [it would be a nice example of group objects in a category].

Topics left out. Nuclear groups, topological vector spaces [these are both important, but I am not sure if I can present them in an elementary way].

Acknowledgements

I am deeply indebted to Reinhardt Winkler and his research group for their hospitality during my visit at Technische Universität Wien, who inspired and encouraged me to organize my thoughts and notes on the topic.

I wish to thank Salvador Hernández, Keith Johnson, and Jeff Egger for the valuable discussions and helpful suggestions that were of great assistance in writing this paper.

Chapter I

The Pontryagin dual

I.A. The evaluation homomorphism

I.1. Pontryagin dual. For Hausdorff topological groups G and H , we denote by $\mathcal{H}(G, H)$ the space of continuous homomorphisms $\varphi: G \rightarrow H$ equipped with the compact-open topology. (For a brief review of the compact-open topology, see Appendix B.) Since the property of being a homomorphism is equationally defined, $\mathcal{H}(G, H)$ is a closed subspace of $\mathcal{C}(G, H)$. Indeed,

$$\mathcal{H}(G, H) = \bigcap_{g_1, g_2 \in G} \{f \in \mathcal{C}(G, H) \mid f(g_1 g_2) = f(g_1) f(g_2)\}, \quad (\text{I.1})$$

and each of the sets in the intersection is closed. Topological groups have natural uniform structures, and the compact-open topology can also be realized as the topology of uniform convergence on compacta: $\varphi_\alpha \rightarrow \varphi$ in $\mathcal{H}(G, H)$ if $\varphi_\alpha|_K$ converges uniformly to $\varphi|_K$ for every compact subset $K \subseteq G$.

For $A \in \text{Ab}(\text{Top})$, a *character* of A is a homomorphism $\chi: A \rightarrow \mathbb{T}$, where $\mathbb{T} := \mathbb{R}/\mathbb{Z}$. If χ is continuous, then it must factor through the maximal Hausdorff quotient A/N_A of A , because $\mathbb{T}$ is Hausdorff (cf. Proposition A.2). The *Pontryagin dual* $\hat{A}$ of A is the group $\mathcal{H}(A/N_A, \mathbb{T})$ of all continuous characters of A equipped with the compact-open topology. Since compact subsets of A/N_A are precisely the images of compact subsets in A , one may view $\hat{A}$ as the group $\mathcal{H}(A, \mathbb{T})$ both algebraically and topologically (cf. Corollary A.4).

We put A_d for the group A equipped with the discrete topology. The space $\mathcal{C}(A_d, \mathbb{T})$ coincides with $\mathbb{T}^{A_d}$, and thus compact. Therefore, its closed subspace $\widehat{A_d}$ is also compact; it carries the topology of *pointwise convergence* on A , in other words, $\chi_\alpha \rightarrow \chi$ if and only if $\chi_\alpha(a) \rightarrow \chi(a)$ for every $a \in A$. Since the compact-open topology is finer than the pointwise convergence one, $\hat{A} \rightarrow \widehat{A_d}$ is continuous.

I.2. Polar. The sets $\Lambda_n = [-\frac{1}{4n}, \frac{1}{4n}]$ form a base at 0 for $\mathbb{T}$, and have the property that if $kx \in \Lambda_1$ for $k = 1, \dots, n$, then $x \in \Lambda_n$. (Here, $[-\frac{1}{4n}, \frac{1}{4n}]$ are identified with their images in $\mathbb{T}$.) For $S \subseteq A$

and $\Phi \subseteq \hat{A}$, their *polar* sets are defined as

$$S^\triangleright = \{\chi \in \hat{A} \mid \chi(S) \subseteq \Lambda_1\}, \quad (\text{I.2})$$

$$\Phi^\triangleleft = \{a \in A \mid \forall \chi \in \Phi, \chi(a) \in \Lambda_1\} = \bigcap_{\chi \in \Phi} \chi^{-1}(\Lambda_1). \quad (\text{I.3})$$

Since $S^\triangleright$ is closed in the topology of pointwise convergence, in particular, it is closed in the compact-open one. On the other hand, $\Phi^\triangleleft$ is closed in A because each $\chi \in \Phi$ is continuous. It is immediately seen that $(\triangleleft, \triangleright)$ is a Galois-connection between subsets of A and $\hat{A}$. Thus, $S \subseteq S'^{\triangleright\triangleleft}$ and $\Phi \subseteq \Phi^{\triangleleft\triangleright}$.

I.3. Lemma. Let $\varphi: A \rightarrow B$ be a morphism in $\text{Ab}(\text{Top})$.

- (a) For $S \subseteq A$, $\hat{\varphi}^{-1}(S^\triangleright) = \varphi(S)^\triangleright$.
- (b) For $\Sigma \subseteq \hat{B}$, $\varphi^{-1}(\Sigma^\triangleleft) = \hat{\varphi}(\Sigma)^\triangleleft$.
- (c) For $S' \subseteq B$, $\hat{\varphi}(S'^\triangleright) \subseteq \varphi^{-1}(S')^\triangleright$.
- (d) For $\Sigma' \subseteq \hat{A}$, $\varphi(\Sigma'^\triangleleft) \subseteq \hat{\varphi}^{-1}(\Sigma')^\triangleleft$.
- (e) For $S \subseteq A$, $\varphi(S^{\triangleright\triangleleft}) \subseteq \varphi(S)^{\triangleright\triangleleft}$.
- (f) For $S' \subseteq B$, $\varphi^{-1}(S')^{\triangleright\triangleleft} \subseteq \varphi^{-1}(S'^{\triangleright\triangleleft})$.

Proof. Since φ is continuous, it induces a continuous homomorphism $\hat{\varphi}: \hat{B} \rightarrow \hat{A}$ defined by $\hat{\varphi}(\chi) = \chi \circ \varphi$.

$$\chi \in \varphi(S)^\triangleright \iff \chi(\varphi(S)) \subseteq \Lambda_1 \iff \chi \circ \varphi \in S^\triangleright \iff \chi \in \hat{\varphi}^{-1}(S^\triangleright), \quad (\text{I.4})$$

and so (a) follows.

$$g \in \varphi^{-1}(\Sigma^\triangleleft) \iff \varphi(g) \in \Sigma^\triangleleft \iff \forall \chi \in \Sigma, \chi(\varphi(g)) \in \Lambda_1 \quad (\text{I.5})$$

$$\iff \forall \psi \in \hat{\varphi}(\Sigma), \psi(g) \in \Lambda_1 \iff g \in \hat{\varphi}(\Sigma)^\triangleleft, \quad (\text{I.6})$$

which shows (b).

(c) Since $\varphi(\varphi^{-1}(S')) \subseteq S'$, one has $S'^\triangleright \subseteq \varphi(\varphi^{-1}(S'))^\triangleright$. By (a) applied to $S = \varphi^{-1}(S')$, $\varphi(\varphi^{-1}(S'))^\triangleright = \hat{\varphi}^{-1}(\varphi^{-1}(S')^\triangleright)$. Thus, $S'^\triangleright \subseteq \hat{\varphi}^{-1}(\varphi^{-1}(S')^\triangleright)$.

(d) Since $\hat{\varphi}(\hat{\varphi}^{-1}(\Sigma')) \subseteq \Sigma'$, one has $\Sigma'^\triangleleft \subseteq \hat{\varphi}(\hat{\varphi}^{-1}(\Sigma'))^\triangleleft$. By (b) applied to $\Sigma = \hat{\varphi}^{-1}(\Sigma')$, $\hat{\varphi}(\hat{\varphi}^{-1}(\Sigma'))^\triangleleft = \varphi^{-1}(\hat{\varphi}^{-1}(\Sigma')^\triangleleft)$. Thus, $\Sigma'^\triangleleft \subseteq \varphi^{-1}(\hat{\varphi}^{-1}(\Sigma')^\triangleleft)$.

(e) follows from (a) and (d), and (f) follows from (b) and (c). $\square$

I.4. Lemma. Let $A \in \text{Ab}(\text{Top})$, $H \leq A$ its subgroup, and let $\pi_H: A \rightarrow A/H$ be the canonical projection. Then:

- (a) $H^\triangleright$ is a subgroup;
- (b) $\hat{\pi}_H: \widehat{A/H} \rightarrow \hat{A}$ is injective, and its image is $H^\triangleright$;
- (c) ([6, 5]) if $S' \subseteq A/H$ and $0 \in S'$, then $\pi_H^{-1}(S')^\triangleright = \hat{\pi}_H(S'^\triangleright)$.

Proof. (a) Since Λ_1 contains only the zero subgroup, $\chi(H) \subseteq \Lambda_1$ holds if and only if $H \subseteq \ker \chi$. Thus, $H^\triangleright$ is the annihilator subgroup of H in $\hat{A}$.

(b) Since π_H is onto, $\hat{\pi}_H$ is injective. Clearly, $\text{Im } \hat{\pi}_H \subseteq H^\triangleright$, so in order to show the converse, let $\chi \in H^\triangleright$. Then $H \subseteq \ker \chi$, and thus χ induces a continuous character $\bar{\chi}: A/H \rightarrow \mathbb{T}$ such that $\hat{\pi}_H(\bar{\chi}) = \bar{\chi} \circ \pi_H = \chi$.

(c) One has $H \subseteq \pi_H^{-1}(S')$ because $0 \in S'$, and thus $\pi_H^{-1}(S')^\triangleright \subseteq H^\triangleright = \text{Im } \hat{\pi}_H$, by (b). Lemma I.3(c) applied to $S' = \pi_H^{-1}(S')$ yields $\hat{\pi}_H^{-1}(\pi_H^{-1}(S')^\triangleright) = \pi_H(\pi_H^{-1}(S'))^\triangleright = S'^\triangleright$. This completes the proof, because $\hat{\pi}_H$ is injective. $\square$

For $X \in \text{Top}$, we denote by $\mathcal{K}(X)$ the collection of compact subsets of X . For $G \in \text{Grp}(\text{Top})$, one puts $\mathcal{N}(G)$ for the collection of neighborhoods of e in G .

I.5. Proposition. Let $A \in \text{Ab}(\text{Top})$.

(a) $\chi \in \text{hom}(A, \mathbb{T})$ is continuous if and only if there is $U \in \mathcal{N}(A)$ such that $\chi(U) \subseteq \Lambda_1$.

(b) The collection $\{K^\triangleright \mid K \subseteq A, K \in \mathcal{K}(A)\}$ is a base at 0 to $\hat{A}$.

(c) $\Sigma \subseteq \hat{A}$ is equicontinuous if and only if there is $U \in \mathcal{N}(A)$ such that $\Sigma \subseteq U^\triangleright$.

Proof. Observe that in order to establish continuity-like properties of homomorphisms of topological groups, it suffices to check them at a single point, and 0 is a convenient choice for that purpose.

(a) Necessity is obvious. So, in order to show sufficiency, let $n \in \mathbb{N}$ and let $U \in \mathcal{N}(A)$ be such that $\chi(U) \subseteq \Lambda_1$. Because of the continuity of addition in A , there is $V \in \mathcal{N}(A)$ such that $\underbrace{V + \dots + V}_{n \text{ times}} \subseteq U$, and in particular, $k\chi(V) \subseteq \Lambda_1$ for every $k = 1, \dots, n$. Therefore, $\chi(V) \subseteq \Lambda_n$, as desired.

(b) Let $C \in \mathcal{K}(A)$, and set $K = C \cup 2C \cup \dots \cup nC$. Then $K \in \mathcal{K}(A)$, and $\chi(C) \subseteq \Lambda_n$ for every $\chi \in K^\triangleright$, because $k\chi(a) \in \Lambda_1$ for every $a \in C$ and $k = 1, \dots, n$. Thus, we obtained $K^\triangleright \subseteq \{\chi \in \hat{A} \mid \chi(C) \subseteq \Lambda_n\}$, as desired.

(c) Because of the homogeneous structure of topological groups, equicontinuity of a family Σ of characters can be checked at a single point. Equicontinuity at 0 means that for every neighborhood Λ_n of 0 there is $V \in \mathcal{N}(A)$ such that $\chi(V) \subseteq \Lambda_n$ for every $\chi \in \Sigma$. By the argument presented in (a), this condition is satisfied for all n if and only if it is satisfied for $n = 1$. Since this condition for $n = 1$ is precisely $\Sigma \subseteq U^\triangleright$, the proof is complete. $\square$

I.6. Evaluation homomorphism. Each $a \in A$ gives rise to a continuous character $\hat{a}$ of $\text{hom}(A, \mathbb{T})$ given by evaluation: $\hat{a}(\psi) = \psi(a)$. In particular, $\hat{a}$ is a continuous character of $\hat{A}$ (whose topology is finer than that of $\text{hom}(A, \mathbb{T})$). Thus, $\hat{a} \in \hat{\hat{A}}$, and the evaluation map

$$\alpha_A: A \longrightarrow \hat{\hat{A}} \tag{I.7}$$

$$a \longmapsto \hat{a} \tag{I.8}$$

is a homomorphism of groups. In the sequel, we present necessary and sufficient conditions for α_A to be continuous and an embedding (cf. Propositions I.7 and I.12), and it will become clear that in general, α_A need not be continuous.

A collection $\mathcal{C}$ of compact sets of a topological space X is a *cobase* if for every compact subset $K \subseteq X$ there is $C \in \mathcal{C}$ such that $K \subseteq C$.

I.7. Proposition. ([28, 2.3]) Let $A \in \text{Ab}(\text{Top})$.

- (a) For every $U \in \mathcal{N}(A)$, $U^\triangleright$ is compact in $\hat{A}$.
- (b) The following statements are equivalent:
 - (i) α_A is continuous;
 - (ii) every compact subset of $\hat{A}$ is equicontinuous;
 - (iii) $\{U^\triangleright \mid U \in \mathcal{N}(A)\}$ is a cobase for $\hat{A}$;
 - (iv) $\{U^{\triangleright\triangleright} \mid U \in \mathcal{N}(A)\}$ is a base for $\hat{A}$ at 0.
- (c) $\ker \alpha_A = \bigcap_{\chi \in \hat{A}} \ker \chi$.

Proof. (a) As noted earlier, $U^\triangleright$ is closed in $\hat{A}$ (it is closed even in $\text{hom}(A, \mathbb{T})$). By Proposition I.5(c), it is also equicontinuous, and therefore it is compact by a standard Arzela-Ascoli type argument.

(b) The equivalence of (ii) and (iii) is an immediate consequence of Proposition I.5(c).

(i) $\Rightarrow$ (ii): Suppose that α_A is continuous, and let $\Sigma \subseteq \hat{A}$ be compact. Then $\Sigma^\triangleright$ is a neighborhood of 0 in $\hat{A}$. Thus, by continuity of α_A , there is $U \in \mathcal{N}(A)$ such that $\alpha_A(U) \subseteq \Sigma^\triangleright$, which is equivalent to $\Sigma \subseteq U^\triangleright$. Therefore, by Proposition I.5(c), Σ is equicontinuous.

(iii) $\Rightarrow$ (iv): If $V \in \mathcal{N}(\hat{A})$, then by Proposition I.5, there is a compact subset $\Phi \subseteq \hat{A}$ such that $\Phi^\triangleright \subseteq V$. Since $\{U^\triangleright \mid U \in \mathcal{N}(A)\}$ is a cobase for $\hat{A}$, there is $U \in \mathcal{N}(A)$ such that $\Phi \subseteq U^\triangleright$. Therefore, $U^{\triangleright\triangleright} \subseteq \Phi^\triangleright \subseteq V$, as desired.

(iv) $\Rightarrow$ (i): If $W \in \mathcal{N}(\hat{A})$, then there is $U \in \mathcal{N}(A)$ such that $U^{\triangleright\triangleright} \subseteq W$, and therefore $\alpha_A(U) \subseteq U^{\triangleright\triangleright} \subseteq W$. Hence, α_A is continuous.

(c) For $a \in A$, one has

$$a \in \ker \alpha_A \iff \alpha_A(a) = 0 \iff \forall \chi \in \hat{A}, (\alpha_A(a))(\chi) = 0 \iff \forall \chi \in \hat{A}, \chi(a) = 0, \quad (\text{I.9})$$

which completes the proof. $\square$

A combination of Propositions I.5(b) and I.7(b) yields:

I.8. Corollary. Let $A \in \text{Ab}(\text{Top})$. The following statements are equivalent:

- (i) $\alpha_{\hat{A}}$ is continuous;
- (ii) $\{K^{\triangleright\triangleright} \mid K \in \mathcal{K}(A)\}$ is a cobase for $\hat{A}$. $\square$

A map $f: X \rightarrow Y$ between Hausdorff spaces is said to be *k-continuous* if the restriction $f|_K$ is continuous for every compact subset K of X . Although α_A need not be continuous (cf. Proposition I.7), its restriction to any compact subset of A is continuous.

I.9. Theorem. Let $A \in \text{Ab}(\text{Haus})$. The evaluation homomorphism α_A is *k-continuous*.

Proof. By Corollary B.3, $e: A \rightarrow \mathcal{C}(\mathcal{C}(A, \mathbb{T}), \mathbb{T})$ is *k-continuous*. Since the image $e(A)$ is contained in $\mathcal{H}(\mathcal{C}(A, \mathbb{T}), \mathbb{T})$, and the map $\mathcal{H}(\mathcal{C}(A, \mathbb{T}), \mathbb{T}) \rightarrow \mathcal{H}(\mathcal{H}(A, \mathbb{T}), \mathbb{T}) = \hat{A}$ (given by restriction to $\mathcal{H}(A, \mathbb{T})$) is continuous, α_A is the composition of a *k-continuous* map with a continuous one. $\square$

I.10. Local quasi-convexity. Let $A \in \text{Ab}(\text{Top})$. Following Banaszczyk, $S \subseteq A$ is *quasi-convex* if $S = S^{\triangleright\triangleleft}$. The group A is *locally quasi-convex* (or briefly, *LQC*) if it admits a base of quasi-convex neighborhoods at 0, that is, if $\{U^{\triangleright\triangleleft} \mid U \in \mathcal{N}(A)\}$ is a base at 0 (cf. [3]).

I.11. Proposition. ([6, 1]) *The group $\hat{A}$ is locally quasi-convex for every $A \in \text{Ab}(\text{Top})$.*

Proof. By Proposition I.5, the collection $\{K^{\triangleright} \mid K \in \mathcal{K}(A)\}$ is a base at 0 at $\hat{A}$. Each member of the collection is quasi-convex, because $K^{\triangleright\triangleright\triangleleft} = K^{\triangleright}$. $\square$

A variant of the next proposition appeared in [2, 6.10] and [9, 4.3], and seems to be a well-known result.

I.12. Proposition. *Let $A \in \text{Ab}(\text{Top})$.*

- (a) *If α_A is an embedding, then A is locally quasi-convex.*
- (b) *If A is locally quasi-convex, then α_A is open onto its image.*
- (c) *If A is locally quasi-convex and Hausdorff, then α_A is injective.*

Proof. First, note that for $S \subseteq A$, one has $\alpha_A(S^{\triangleright\triangleleft}) = S^{\triangleright\triangleright} \cap \alpha_A(A)$. If $U \in \mathcal{N}(A)$, then $U^{\triangleright}$ is compact in $\hat{A}$ (cf. Proposition I.7(a)), and so $U^{\triangleright\triangleright}$ is open in $\hat{A}$ by Proposition I.5(b). Therefore, $\alpha_A(U^{\triangleright\triangleleft})$ is open in $\alpha_A(A)$.

(a) Since α_A is an embedding, $\alpha_A(V)$ is open in $\alpha_A(A)$ for every $V \in \mathcal{N}(A)$. Thus, by Proposition I.7(b)(iv), there is $U \in \mathcal{N}(A)$ such that $\alpha_A(U^{\triangleright\triangleleft}) = U^{\triangleright\triangleright} \cap \alpha_A(A) \subseteq \alpha_A(V)$, which implies $U^{\triangleright\triangleleft} \subseteq V$, because α_A is injective.

(b) Let $V \in \mathcal{N}(A)$, and using LQC pick $U \in \mathcal{N}(A)$ such that $U^{\triangleright\triangleleft} \subseteq V$. Then one has $\alpha_A(U^{\triangleright\triangleleft}) \subseteq \alpha_A(V)$, and $\alpha_A(U^{\triangleright\triangleleft})$ is open in $\alpha_A(A)$.

(c) Let $a \in A$ be a non-zero element. Since A is Hausdorff and LQC, there exists $U \in \mathcal{N}(A)$ such that $a \notin U^{\triangleright\triangleleft}$. Thus, there is $\chi \in U^{\triangleright}$ such that $(\alpha_A(a))(\chi) = \chi(a) \notin \Lambda_1$. Therefore, $\alpha_A(a) \neq 0$. Hence, α_A is injective. $\square$

I.13. Remarks. (1) In Proposition I.12(b), the map α_A need not be an embedding, because it is not necessarily continuous or injective.

(2) Following Neumann and Wigner, whenever α_A is injective, A is said to be *maximally almost-periodic* (or briefly, *MAP*; cf. [27]).

(3) For every $A \in \text{Ab}(\text{Top})$, $\alpha_{\hat{A}}$ is injective. Indeed, if $\alpha_{\hat{A}}(\chi) = 0$, then for every $\xi \in \hat{\hat{A}}$, $\xi(\chi) = (\alpha_{\hat{A}}(\chi))(\xi) = 0$. Thus, for $a \in A$ and $\xi = \alpha_A(a)$, one obtains $\chi(a) = (\alpha_A(a))(\chi) = 0$. Therefore, $\chi = 0$, as desired.

I.14. Corollary. *Let $A \in \text{Ab}(\text{Haus})$ be such that α_A is continuous. Then α_A is an embedding if and only if A is locally quasi-convex.* $\square$

We denote by LQC the full subcategory of $\text{Ab}(\text{Top})$ formed by the locally quasi-convex groups, and present a functorial method of “turning” every group into an LQC group.

I.15. Theorem. *LQC is an epireflective subcategory of $\text{Ab}(\text{Top})$, and the reflection A^Δ is given by equipping the group of A with a new group topology whose base at 0 is $\{U^{\triangleright\triangleleft} \mid U \in \mathcal{N}(A)\}$.*

In less categorical language, Theorem I.15 states that for every $A \in \text{Ab}(\text{Top})$:

- (1) A^Δ is a topological group;
- (2) there is a continuous surjective homomorphism $\nu_A: A \rightarrow A^\Delta$ that is natural in A ;
- (3) $\varphi^\Delta: A^\Delta \rightarrow B^\Delta$ is continuous whenever $\varphi: A \rightarrow B$ is a continuous homomorphism;
- (4) Every continuous homomorphism $\gamma: A \rightarrow C$ into an LQC group C factors uniquely through ν_A , that is, there is a unique $\gamma': A^\Delta \rightarrow C$ such that $\gamma = \gamma' \circ \nu_A$.

Proof. (1) Each set of the form $U^{\triangleright\triangleleft}$ is symmetric (i.e., $-U^{\triangleright\triangleleft} = U^{\triangleright\triangleleft}$), and it is easily seen that $(U \cap V)^{\triangleright\triangleleft} \subseteq U^{\triangleright\triangleleft} \cap V^{\triangleright\triangleleft}$. Let $U \in \mathcal{N}(A)$, and let $V \in \mathcal{N}(A)$ be such that $V + V \subseteq U$. Then $V^{\triangleright\triangleleft} + V^{\triangleright\triangleleft} \subseteq (V + V)^{\triangleright\triangleleft} \subseteq U^{\triangleright\triangleleft}$. Therefore, the proposed candidate for $\mathcal{N}(A^\Delta)$ defines a group topology on the underlying group of A , as desired.

(2) The topology of A^Δ is coarser than that of A , so the identity homomorphism $A \rightarrow A^\Delta$ is continuous (and it is obviously natural in A).

(3) Let $U^{\triangleright\triangleleft} \in \mathcal{N}(B^\Delta)$. By Lemma I.3(f), $\varphi^{-1}(U^{\triangleright\triangleleft}) \supseteq \varphi^{-1}(U)^{\triangleright\triangleleft}$. Since φ is continuous, $\varphi^{-1}(U) \in \mathcal{N}(A)$, and therefore $\varphi^{-1}(U)^{\triangleright\triangleleft} \in \mathcal{N}(A^\Delta)$. Hence, φ^Δ is continuous.

(4) Since A and A^Δ have the same underlying set, uniqueness of γ' is clear. If $C \in \text{LQC}$, then $C = C^\Delta$, and therefore $\gamma' = \gamma^\Delta$, by (3). $\square$

I.16. Corollary. *The limit of a family of locally quasi-convex groups formed in $\text{Ab}(\text{Top})$ is locally quasi-convex, and coincides with the limit formed in LQC. In particular, LQC is closed under the formation of arbitrary products and subgroups.*

Proof. The first statement is a well-known category theoretical property of reflective subcategories (cf. [24, IV.3, V.5]), and it implies the second one, as products are limits. For the third statement, let A be an LQC group and $H \leq A$ be its subgroup. The reflection $(A/H)^\Delta$ has the same underlying group as A/H , and therefore

$$H = \text{Eq}(A \xrightarrow[\quad]{\pi_H^\Delta} (A/H)^\Delta), \quad (\text{I.10})$$

where $\pi_H: A \rightarrow A/H$ is the canonical projection. Since equalizers are limits, this completes the proof. $\square$

I.17. Corollary. *For every $A \in \text{Ab}(\text{Top})$, the underlying groups of $\hat{A}$ and $\widehat{A^\Delta}$ coincide.*

Proof. By Proposition I.11 the group $\hat{\mathbb{Z}} = \mathbb{T}$ is LQC. Thus, by Theorem I.15, every continuous character $\chi: A \rightarrow \mathbb{T}$ gives rise to a continuous $\chi^\Delta: A^\Delta \rightarrow \mathbb{T}$. Therefore, $\hat{A} \subseteq \widehat{A^\Delta}$ as sets. The reverse inclusion is obvious, because A^Δ carries a coarser topology than A . $\square$

I.18. Proposition. *Let $A \in \text{Ab}(\text{Top})$ such that α_A is continuous. Then:*

- (a) $\hat{A}$ and $\widehat{A^\Delta}$ have the same compact subsets.
- (b) A^Δ coincides with the group A equipped with the initial topology induced by α_A ;
- (c) $N_{A^\Delta} = \ker \alpha_{A^\Delta} = \ker \alpha_A$;
- (d) α_{A^Δ} is continuous and open onto its image;
- (e) $\hat{\nu}_A: \hat{A} \rightarrow \widehat{A^\Delta}$ is an embedding;

Proof. Since $\hat{\nu}_A: \widehat{A^\Delta} \rightarrow \hat{A}$ is continuous, every compact subset Φ of $\widehat{A^\Delta}$ is also compact in $\hat{A}$, and thus, by Proposition I.7(b), there exists $U \in \mathcal{N}(A)$ such that $\Phi \subseteq U^\triangleright$.

(a) Suppose that $\Sigma \subseteq \hat{A}$ is compact. Then, by Proposition I.7(b), $\Sigma \subseteq U^\triangleright$ for some $U \in \mathcal{N}(A)$. By Proposition I.7(a), $U^\triangleright = (U^{\triangleright\triangleleft})^\triangleright$ is compact in $\widehat{A^\Delta}$. Since $\Sigma = \hat{\nu}_A^{-1}(\Sigma)$ is a closed subset of $U^\triangleright$, Σ is compact in $\widehat{A^\Delta}$.

(b) By Proposition I.7(b), $\{U^{\triangleright\triangleright} \mid U \in \mathcal{N}(A)\}$ is a base for $\hat{A}$ at 0, because α_A is continuous. The statements follows from $\alpha_A^{-1}(U^{\triangleright\triangleright}) = U^{\triangleright\triangleleft}$.

(c) The first equality follows from (b) and Proposition A.2, because $\hat{A}$ is Hausdorff. For the second equality, observe that by Proposition I.7(c), $\ker \alpha_A = \bigcap_{\chi \in \hat{A}} \ker \chi$, and by (a), A and A^Δ have the same continuous characters.

(d) By (a), polars of quasi-convex neighborhoods form a cobase to $\widehat{A^\Delta}$, and thus α_{A^Δ} is continuous, by Proposition I.7(b). Therefore, the statement follows by Proposition I.12(b).

(e) By Proposition I.7(b), the collections $\{U^{\triangleright\triangleright} \mid U \in \mathcal{N}(A)\}$ and $\{V^{\triangleright\triangleright} \mid V \in \mathcal{N}(A^\Delta)\}$ are bases at 0 to $\hat{A}$ and $\widehat{A^\Delta}$, respectively. Since every $V \in \mathcal{N}(A^\Delta)$ has the form of $U^{\triangleright\triangleleft}$, this completes the proof. $\square$

I.19. Theorem. Let $A \in \text{Ab}(\text{Top})$, and $D \leq A$ be a dense subgroup. Then:

- (a) the underlying groups of $\hat{D}$ and $\hat{A}$ coincide;
- (b) if α_D is continuous, $\hat{D}$ and $\hat{A}$ have the same compact subsets, and α_A is continuous;
- (c) if D is locally quasi-convex, then so is A .

Proof. (a) The inclusion $\iota_D: D \rightarrow A$ induces a continuous homomorphism $\hat{\iota}_D: \hat{A} \rightarrow \hat{D}$, which is injective, because D is dense in A . In order to show that $\hat{\iota}_D$ is surjective, let $\chi \in \hat{D}$. For each $a \in A$, there is a net $\{x_\alpha\} \subseteq D$ such that $x_\alpha \rightarrow a$. Thus, $x_\alpha - x_\beta \rightarrow 0$, and so $\chi(x_\alpha) - \chi(x_\beta) \rightarrow 0$. In other words, $\{\chi(x_\alpha)\}$ is a Cauchy net in the complete group $\mathbb{T}$, and therefore $\lim \chi(x_\alpha)$ exists. Set $\bar{\chi}(a) = \lim \chi(x_\alpha)$; it is easily seen that $\bar{\chi}$ is well-defined and continuous. Hence, $\hat{\iota}_D(\bar{\chi}) = \chi$, as desired.

(b) It suffices to show that $\hat{\iota}_D^{-1}(\Phi)$ is compact in $\hat{A}$ for every compact subset $\Phi \subseteq \hat{D}$, because $\hat{\iota}_D$ is continuous. By Proposition I.7(b), since α_D is continuous, Φ is equicontinuous on the dense subgroup D , which means that there is $U \in \mathcal{N}(A)$ such $\Phi \subseteq (U \cap D)^\triangleright$ (cf. Proposition I.5(b)). It follows from the density of D that $\text{Int } U \subseteq \overline{D \cap U}$, and therefore

$$\Phi \subseteq (U \cap D)^\triangleright = (\overline{U \cap D})^\triangleright \subseteq (\text{Int } U)^\triangleright. \quad (\text{I.11})$$

Hence, $\hat{\iota}_D^{-1}(\Phi)$ is a closed subset of the compact subset $(\text{Int } U)^\triangleright$ of $\hat{A}$ (cf. Proposition I.5(a)). Since $\text{Int } U \in \mathcal{N}(A)$, this also shows that every compact subset of $\hat{A}$ is contained in $U^\triangleright$ for some $U \in \mathcal{N}(A)$, which means, by Proposition I.7(b), that α_A is continuous.

(c) Let $V \in \mathcal{N}(A)$, and pick $V_1 \in \mathcal{N}(A)$ such that $V_1 - V_1 \subseteq V$, so $\bar{V}_1 \subseteq \text{Int } V$ (cf. Proposition A.1(a)). There exists $W_1 \in \mathcal{N}(D)$ such that $W_1^{\triangleright\triangleleft} \cap D \subseteq V_1 \cap D$, because $V_1 \cap D \in \mathcal{N}(D)$ and D is LQC. (For $\Sigma \subseteq \hat{D}$, $\Sigma^{\triangleleft}$ in D is $\hat{\iota}_D^{-1}(\Sigma)^{\triangleleft} \cap D$, because $\hat{D} = \hat{A}$ as sets.) There is $W_2 \in \mathcal{N}(A)$ such that $W_1 = W_2 \cap D$, and since D is dense, one has $W_1^\triangleright = W_2^\triangleright$. Thus, $W_1^{\triangleright\triangleleft} = W_2^{\triangleright\triangleleft}$, and therefore

$$\text{Int}(W_2^{\triangleright\triangleleft}) \subseteq \overline{W_2^{\triangleright\triangleleft} \cap D} = \overline{W_1^{\triangleright\triangleleft} \cap D} \subseteq \overline{V_1 \cap D} = \bar{V}_1 \subseteq V. \quad (\text{I.12})$$

For $U \in \mathcal{N}(A)$ such that $U + U \subseteq W_2$, one has $U^{\triangleright\triangleleft} + U^{\triangleright\triangleleft} \subseteq W_2^{\triangleright\triangleleft}$, and so $U^{\triangleright\triangleleft} \subseteq \text{Int}(W_2^{\triangleright\triangleleft})$ (cf. Proposition A.1(a)). Hence, we found $U \in \mathcal{N}(A)$ such that $U^{\triangleright\triangleleft} \subseteq V$, as desired. $\square$

I.20. Corollary. *The completion of every locally quasi-convex group is locally quasi-convex.* $\square$

I.21. Precompactness. A subset $S \subseteq G$ of $G \in \text{Grp}(\text{Top})$ is said to be *precompact* if for every $U \in \mathcal{N}(A)$ there is a finite subset $F \subseteq G$ such that $S \subseteq FU$. Similarly to compactness, if $S, S' \subseteq G$ are precompact and $S_1 \subseteq S$, then so are $S_1, \bar{S}, S + S'$, and $\varphi(S)$ for every continuous homomorphism $\varphi: G \rightarrow H$. Our interest in this property arises from the following theorem on uniform spaces (cf. [17] and [13, 8.3.16]):

I.22. Theorem. *A uniform space $(X, \mathcal{U})$ is compact if and only if it is complete and precompact.*

I.23. Proposition. *Let $A \in \text{Ab}(\text{Haus})$ be a locally quasi-convex group. Then:*

- (a) $\alpha_A^{-1}(E)$ is precompact in A for every equicontinuous subset $E \subseteq \hat{\hat{A}}$;
- (b) $K^{\triangleright\triangleleft}$ is precompact in A for every compact subset $K \subseteq A$;
- (c) if $\alpha_{\hat{A}}$ is continuous, then $\alpha_A^{-1}(F)$ is precompact for every compact $F \subseteq \hat{\hat{A}}$.

Proof. First, we note that by Proposition I.12, $\alpha_A^{-1}: \alpha_A(A) \rightarrow A$ is continuous.

(a) By Proposition I.5(c), there is $V \in \mathcal{N}(\hat{A})$ such that $E \subseteq V^{\triangleright}$. By Proposition I.7(a), $V^{\triangleright}$ is compact in $\hat{\hat{A}}$, and so $V^{\triangleright} \cap \alpha_A(A)$ is precompact. Thus, the continuous homomorphic image $\alpha_A^{-1}(V^{\triangleright} \cap \alpha_A(A))$ is also precompact, and contains $\alpha_A^{-1}(E)$. Therefore, $\alpha_A^{-1}(E)$ is precompact.

(b) The set $K^{\triangleright}$ is open in $\hat{A}$, and thus $K^{\triangleright\triangleright}$ is compact and equicontinuous in $\hat{\hat{A}}$ (cf. Propositions I.7(a) and I.5(c)). Hence, by (a), $K^{\triangleright\triangleleft} = \alpha_A^{-1}(K^{\triangleright\triangleright})$ is precompact in A .

(c) By Propositions I.8, in this case, $\{K^{\triangleright\triangleright} \mid K \in \mathcal{K}(A)\}$ is a cobase for $\hat{\hat{A}}$, and so the statement follows from (b). $\square$

I.24. Quasi-convex compactness. It is a well-known that the closed convex hull of a weakly compact subset of a Banach space is weakly compact. In fact, the property of preservation of compactness under formation of closed convex hull characterizes completeness in the category of metrizable locally convex spaces (cf. [29, 2.4]). Motivated by this, one says that $A \in \text{Ab}(\text{Top})$ has the *quasi-convex compactness property* (or briefly, A is *QCP*) if for every compact subset $K \subseteq A$, the quasi-convex hull $K^{\triangleright\triangleleft}$ (of K in A) is compact.

I.25. Proposition. *Let $A \in \text{Ab}(\text{Top})$ be such that α_A is continuous. Then $\hat{A}$ has the quasi-convex compactness property.*

Proof. Let $\Phi \subseteq \hat{A}$ be compact. By Proposition I.7(b), there is $U \in \mathcal{N}(A)$ such that $\Phi \subseteq U^{\triangleright}$, and so $\Phi^{\triangleright\triangleleft} \subseteq U^{\triangleright\triangleright\triangleleft} = U^{\triangleright}$. Therefore, the closed set $\Phi^{\triangleright\triangleleft}$ is a subset of the compact set $U^{\triangleright}$ (cf. Proposition I.7(a)). Hence, the result follows. $\square$

I.26. Proposition. *Let $A \in \text{Ab}(\text{Haus})$ be a locally quasi-convex group. If*

- (a) A is complete, or
- (b) α_A is surjective,

then A has the quasi-convex compactness property.

Proof. By Proposition I.12, $\alpha_A^{-1}: \alpha_A(A) \rightarrow A$ is continuous.

(a) By Proposition I.23, $K^{\triangleright\triangleleft}$ is precompact in A , and it, being a closed subspace, is complete. Therefore, by Theorem I.22, it is compact.

(b) Since $K^{\triangleright}$ is open in $\hat{A}$, $K^{\triangleright\triangleright}$ is compact in $\hat{A}$, and therefore $K^{\triangleright\triangleleft} = \alpha_A^{-1}(K^{\triangleright\triangleright})$ is compact, because α_A^{-1} is continuous. $\square$

I.27. Proposition. *Let $A \in \text{Ab}(\text{Haus})$ be such that α_A and $\alpha_{\hat{A}}$ are continuous. The following statements are equivalent:*

- (i) $\alpha_A^{-1}(F)$ is compact for every compact subset $F \subseteq \hat{A}$;
- (ii) A has the quasi-convex compactness property.

Proof. (i) $\Rightarrow$ (ii) is immediate, because $K^{\triangleright\triangleright}$ is compact in $\hat{A}$ and $K^{\triangleright\triangleleft} = \alpha_A^{-1}(K^{\triangleright\triangleright})$.

(ii) $\Rightarrow$ (i): Let $F \subseteq \hat{A}$ be compact. Since F is closed and α_A is continuous, $\alpha_A^{-1}(F)$ is also closed in A . By Corollary I.8, there is a compact subset $K \subseteq A$ such that $F \subseteq K^{\triangleright\triangleright}$ (because $\alpha_{\hat{A}}$ is continuous). Thus, $\alpha_A^{-1}(F) \subseteq \alpha_A^{-1}(K^{\triangleright\triangleright}) = K^{\triangleright\triangleleft}$, and $K^{\triangleright\triangleleft}$ is compact because A is QCP. Therefore, $\alpha_A^{-1}(F)$, being the closed subset of a compact set, is compact. $\square$

I.B. Special groups and subgroups

I.28. Metrizable groups. Every Hausdorff topological group is completely regular (nice reference??), so by Tychonoff's metrization theorem, if a Hausdorff group is second countable, then it is metrizable. But for topological groups, second countability is more than necessary in order to warrant metrizability. Theorem I.29 below (originally proved by Kakutani and Birkhoff in 1936) can also be obtained from well-known results on uniform spaces (cf. [13, 8.1.10, 8.1.21]).

I.29. Theorem. *Every first countable Hausdorff topological group is metrizable.* $\square$

We turn to abelian metrizable groups. Recall that a Hausdorff space X is a k -space if $F \subseteq X$ is closed in X whenever $F \cap K$ is closed for every $K \in \mathcal{K}(X)$. (For details, see Appendix C.)

I.30. Theorem. ([10, Theorem 1]) *Let $A \in \text{Ab}(\text{Met})$. Then $\hat{A}$ is a k -space.*

Although Theorem I.30 has a non-commutative generalization (cf. [23], [22, 3.4]), we provide here the original proof by Chasco, enriched with a few words of explanation.

Proof. In order to show that $\hat{A}$ is a k -space, let $\Phi \subseteq \hat{A}$ be such that $\Phi \cap \Xi$ is closed in Ξ for every compact subset $\Xi \subseteq \hat{A}$. We show that for every $\zeta \notin \Phi$, there exists a compact subset $K \subseteq A$ such that $(K^{\triangleright} + \zeta) \cap \Phi = \emptyset$, and so Φ is closed in $\hat{A}$. Without loss of generality, we may assume that $\zeta = 0$.

Since A is metrizable, 0 has a decreasing countable base $\{U_n\}$, and we set $U_0 = A$. Our aim is to construct inductively a family $\{F_n\}_{n=0}^{\infty}$ of finite subsets of A such that for every $n \in \mathbb{N}$,

$$F_n \subseteq U_n, \tag{I.13}$$

$$\bigcap_{k=0}^n F_k^{\triangleright} \cap U_{n+1}^{\triangleright} \cap \Phi = \emptyset. \tag{I.14}$$

By Proposition I.7(a), $U_1^\triangleright$ is compact in $\hat{A}$, and thus by our assumption, $U_1^\triangleright \cap \Phi$ is closed in $U_1^\triangleright$. Thus, $U_1^\triangleright \cap \Phi$ is compact in $\hat{A}$, and so it is compact in the pointwise topology carried by $\text{hom}(A, \mathbb{T})$ (which is coarser than the compact-open one). In particular, $U_1^\triangleright \cap \Phi$ is closed in the pointwise topology. A basic neighborhood of 0 in the pointwise topology has the form $F^\triangleright$, where $F \subseteq A$ is finite, so $0 \notin U_1^\triangleright \cap \Phi$ implies that there exists a finite subset $F_0 \subseteq A = U_0$ such that $F_0^\triangleright \cap U_1^\triangleright \cap \Phi = \emptyset$. This completes the proof for $n = 0$.

Suppose that we have already constructed $F_0, \dots, F_{n-1}$ such that (I.13) and (I.14) hold. For each $x \in U_n$, put

$$\Delta_x = \bigcap_{k=0}^{n-1} F_k^\triangleright \cap \{x\}^\triangleright \cap U_{n+1}^\triangleright \cap \Phi. \quad (\text{I.15})$$

Since $U_n \supseteq U_{n+1}$, one has $U_n^\triangleright \subseteq U_{n+1}^\triangleright$, and thus, by the inductive hypothesis,

$$\bigcap_{x \in U_n} \Delta_x = \bigcap_{k=0}^{n-1} F_k^\triangleright \cap U_n^\triangleright \cap U_{n+1}^\triangleright \cap \Phi = \bigcap_{k=0}^{n-1} F_k^\triangleright \cap U_n^\triangleright \cap \Phi = \emptyset. \quad (\text{I.16})$$

The Δ_x are closed subsets of the compact space $U_{n+1}^\triangleright$, and their intersection is empty. Therefore, there must be a finite subset $F_n \subseteq U_n$ such that $\bigcap_{x \in F_n} \Delta_x = \emptyset$. Hence,

$$\bigcap_{x \in F_n} \Delta_x = \bigcap_{k=0}^{n-1} F_k^\triangleright \cap F_n^\triangleright \cap U_{n+1}^\triangleright \cap \Phi = \emptyset, \quad (\text{I.17})$$

as desired. Set $K = \bigcup_{n=0}^{\infty} F_n \cup \{0\}$. It follows from (I.13) that $K \setminus U_n$ is finite for every $n \in \mathbb{N}$, and thus K is sequentially compact (because every sequence without a constant subsequence must converge to $0 \in K$). Therefore, K is compact, because A is metrizable. By the construction of K , $K^\triangleright \cap U_n^\triangleright \cap \Phi = \emptyset$ for every $n \in \mathbb{N}$. By Proposition I.5(a), $\hat{A} = \bigcup_{n=1}^{\infty} U_n^\triangleright$, and therefore $K^\triangleright \cap \Phi = \emptyset$, as desired. $\square$

I.31. Corollary. *Let $A \in \text{Ab}(\text{Met})$. Then:*

- (a) α_A is continuous;
- (b) $\alpha_{\hat{A}}$ is continuous;
- (c) ([10, Corollary 1]) $\hat{\hat{A}}$ is complete and metrizable.

Proof. (a) Since A is metrizable, its topology is determined by convergent sequences $x_n \longrightarrow x_0$. For such a sequence, $\{x_n \mid n \in \mathbb{N}\} \cup \{x_0\}$ is compact, and thus A is a k -space. Therefore, by Theorem I.9(b), α_A is continuous.

(b) By Theorem I.30, $\hat{A}$ is a k -space, and so, by Theorem I.9(b), $\alpha_{\hat{A}}$ is continuous.

(c) By Proposition I.7(b), $\{U^{\triangleright\triangleright} \mid U \in \mathcal{N}(A)\}$ is a base for $\hat{\hat{A}}$ at 0, because α_A is continuous. Thus, $\hat{\hat{A}}$ is first-countable, because A is so. (Recall that for topological groups, being first-countable is equivalent to being metrizable.) By Theorem I.30, $\hat{A}$ is a k -space, and thus $\mathcal{C}(\hat{A}, \mathbb{T})$ is

complete, because $\mathbb{T}$ is so (cf. [20, 7.12]). Since $\hat{\hat{A}}$ is a closed subspace of $\mathcal{C}(\hat{A}, \mathbb{T})$, this completes the proof. $\square$

A combination of Proposition I.27 and Corollary I.31 yields:

I.32. Corollary. *Let $A \in \text{Ab}(\text{Met})$. Then A has the quasi-convex compactness property if and only if $\alpha_A^{-1}(F)$ is compact in A for every compact $F \subseteq \hat{\hat{A}}$.* $\square$

I.33. Corollary. ([10, Theorem 2], [22, 3.7]) *Let $A \in \text{Ab}(\text{Met})$, and let $D \leq A$ be its dense subgroup. Then $\hat{D} = \hat{A}$ as topological groups.*

Proof. By Theorem I.19, $\hat{D}$ and $\hat{A}$ have the same underlying groups and compact subsets. On the other hand, by Theorem I.30, both $\hat{D}$ and $\hat{A}$ are k -spaces, because D and A are metrizable. Therefore, $\hat{A}$ and $\hat{D}$ have the same topology, because the topology of a k -space is determined by its compact subsets. $\square$

I.34. Theorem. *Let $A \in \text{Ab}(\text{Met})$. The following statements are equivalent:*

- (i) *A is locally quasi-complete and complete;*
- (ii) *α_A is a closed embedding;*
- (iii) *A has the quasi-convex compactness property and α_A is injective.*

Ostling and Wilansky showed that a locally convex metrizable vector space is complete if and only if the absolutely convex closure of compact subset is compact (cf. [29, 2.4]). Hernández proved a far reaching generalization of this result for metrizable groups, namely, that a metrizable LQC group is complete if and only if it has QCP (cf. [15, Theorem 2]). The same result also appears in a paper by Bruguera and Martín-Peinador, who used Corollary I.33 in order to simplify the proof (cf. [8, 9]). It escaped the attention of these authors that if α_A is injective, then QCP is sufficient to warrant completeness, so the first half of the proof of (iii) $\Rightarrow$ (i) appears to be new, while its second half uses the ideas of Bruguera and Martín-Peinador.

Proof. (i) $\Rightarrow$ (ii): Since A is LQC (and metrizable, so Hausdorff), α_A is an embedding by Proposition I.12. Furthermore, if $S \subseteq A$ is closed in A , then S is complete (because A is complete), and thus $\alpha_A(S)$ is complete. Therefore, $\alpha_A(S)$ is closed in $\hat{\hat{A}}$, as desired.

(ii) $\Rightarrow$ (iii): Clearly, α_A is injective. Let $K \subseteq A$ be compact. Then $K^{\triangleright\triangleright}$ is compact in $\hat{\hat{A}}$, and since $\alpha_A(A)$ is closed, $K^{\triangleright\triangleright} \cap \alpha_A(A)$ is also compact. Therefore, $K^{\triangleright\triangleleft} = \alpha_A^{-1}(K^{\triangleright\triangleright} \cap \alpha_A(A))$ is compact too.

(iii) $\Rightarrow$ (i): Since α_A is injective, $\alpha_A^{-1}: \alpha_A(A) \rightarrow A$ is well-defined. By Corollary I.32, QCP implies that $\alpha_A^{-1}(F)$ is compact for every compact subset $F \subseteq \alpha_A(A)$. So, if $K \subseteq \alpha_A^{-1}(F)$ is closed, then K is also compact, and thus $(\alpha_A^{-1})^{-1}(K) = \alpha_A(K)$ is a compact subset of F , because α_A is continuous. Therefore, $(\alpha_A^{-1})^{-1}(K)$ is closed, and hence, α_A^{-1} is k -continuous. By Corollary I.31(c), $\hat{\hat{A}}$ is metrizable, and so is its subspace $\alpha_A(A)$. In particular, $\alpha_A(A)$ is a k -space, and therefore α_A^{-1} is continuous. This shows that α_A is an embedding, and hence, by Proposition I.12(a), A is LQC.

In order to show that A is complete, let B be the completion of A . By Corollary I.33, $\hat{A} = \hat{B}$ (as topological groups), because A is a dense subgroup of the metrizable group B . Let $\{a_n\}$ be a

Cauchy-sequence in A . Then $a_n \longrightarrow b$ for some $b \in B$, and $K_1 = \{a_n \mid n \in \mathbb{N}\} \cup \{b\}$ is compact in B . So, $K_1^\triangleright \in \mathcal{N}(\hat{B}) = \mathcal{N}(\hat{A})$, and by Proposition I.5(b), there is a compact subset $K \subseteq A$ such that $K^\triangleright \subseteq K_1^\triangleright$. Thus, $\{a_n \mid n \in \mathbb{N}\} \subseteq K_1^{\triangleright\triangleleft} \subseteq K^{\triangleright\triangleleft}$ (where $(-)^{\triangleleft}$ is taken with respect to A). Since A has QCP, $K^{\triangleright\triangleleft}$ is compact, and so sequentially compact (because A is metrizable). Therefore, $\{a_n\}$ has a convergent subsequence in $K^{\triangleright\triangleleft}$, which shows that $b \in K^{\triangleright\triangleleft} \subseteq A$. Hence, A is complete, as desired. $\square$

I.35. Compact and discrete groups. Let $A \in \text{Ab}(\text{Top})$. If A is compact, then by Proposition I.5(b), $\{0\} = A^\triangleright \in \mathcal{N}(\hat{A})$. On the other hand, if A is discrete, then $\{0\} \in \mathcal{N}(A)$, and so by Proposition I.7(a), $\hat{A} = \{0\}^\triangleright$ is compact. Repeating these arguments for $\hat{A}$ yields:

I.36. Lemma. Let $A \in \text{Ab}(\text{Top})$.

- (a) If A is compact, then $\hat{A}$ is discrete.
- (b) If A is discrete, then $\hat{A}$ is compact.
- (c) If A is compact, then so is $\hat{\hat{A}}$.
- (d) If A is discrete, then so is $\hat{\hat{A}}$.

$\square$

I.37. Theorem. Let $A \in \text{Ab}(\text{Top})$. If A is

- (a) discrete, or
- (b) compact Hausdorff,

then α_A is an isomorphism of topological groups.

Proof. (a) Let $a \in A$ be a non-zero element, and let $r \in \mathbb{T}$ be such that $o(a) = o(r)$. This defines a group homomorphism $\chi_a: \langle a \rangle \rightarrow \mathbb{T}$ such that $\chi_a(a) \neq 0$. Since $\mathbb{T}$ is an injective abelian group, χ_a extends to $\bar{\chi}_a: A \rightarrow \mathbb{T}$, and $\bar{\chi}_a \in \hat{A}$, because A is discrete. Thus, $\alpha_a(\bar{\chi}_a) = \bar{\chi}_a(a) \neq 0$, which shows that α_A is injective. By Lemma I.36(d), $\hat{\hat{A}}$ is discrete, and thus α_A is an embedding.

Suppose that A is finitely generated. Then it decomposes into the direct sum of cyclic groups $A = \langle a_1 \rangle \oplus \cdots \oplus \langle a_n \rangle$. Finite cyclic groups are self-dual, and $\hat{\mathbb{Z}} = \mathbb{T}$ and $\hat{\mathbb{T}} = \mathbb{Z}$. Thus, surjectivity of α_A follows, because the Pontryagin dual is an additive functor. (More details???)

In the general case, let $\xi \in \hat{\hat{A}}$. By Proposition I.5(a), there is $W \in \mathcal{N}(\hat{A})$ such that $\xi(W) \subseteq \Lambda_1$, and W can be chosen to have the form $\widehat{F^\triangleright}$ where $F \subseteq A$ is finite. Put $B = \langle F \rangle$. By Lemma I.4, $B^\triangleright$ is a subgroup of $\hat{A}$, and $B^{\triangleright\triangleright} \cong \widehat{\hat{A}/B^\triangleright}$. Since $\mathbb{T}$ is an injective abelian group, every homomorphism $\chi: B \rightarrow \mathbb{T}$ extends to a homomorphism $\bar{\chi}: A \rightarrow \mathbb{T}$, and thus the dual $\hat{\iota}_B: \hat{A} \rightarrow \hat{B}$ of the inclusion $\iota_B: B \rightarrow A$ is surjective. The map $\hat{\iota}_B$ is also closed, because $\hat{A}$ is compact. Therefore, $\hat{B} \cong \hat{A}/B^\triangleright$, and hence $B^{\triangleright\triangleright} \cong \widehat{\hat{A}/B^\triangleright} \cong \hat{\hat{B}}$. Since $F \subseteq B$, one has $\xi(B^\triangleright) \subseteq \xi(F^\triangleright) \subseteq \Lambda_1$, and so $\xi \in B^{\triangleright\triangleright} \cong \hat{\hat{B}}$. The group B is finitely generated, and thus by what we have shown so far, there is $b \in B$ such that $\xi = \alpha_B(b)$. Therefore, $\xi = \alpha_A(b)$.

(b) Since A is compact Hausdorff, it is a k -space, and so α_A is continuous (cf. Theorem I.9). By the Peter-Weyl theorem, for every non-zero $a \in A$, there is $\chi \in \hat{A}$ such that $\chi(a) \neq 0$ (cf. [30, Thm. 32]). Thus, α_A is injective, and therefore it is a closed embedding (because A is compact). In order to show that α_A is surjective, assume the contrary, that is, assume that $\alpha_A(A)$ is a proper closed subgroup of $\hat{\hat{A}}$. By Lemma I.36(c), $\hat{\hat{A}}$ is compact, and thus so is the quotient

$\hat{\hat{A}}/\alpha_A(A)$. By the Peter-Weyl theorem applied to $\hat{\hat{A}}/\alpha_A(A)$, there exists a non-zero continuous character $\psi: \hat{\hat{A}}/\alpha_A(A) \rightarrow \mathbb{T}$, which induces $\zeta \in \hat{\hat{A}}$ such that $\zeta(\alpha_A(A)) = \{0\}$ and $\zeta \neq 0$. Since $\hat{A}$ is discrete (cf. Lemma I.36(a)), by (a), $\alpha_{\hat{A}}$ is an isomorphism. In particular, there is a non-zero $\chi \in \hat{A}$ such that $\alpha_{\hat{A}}(\chi) = \zeta$. Therefore, for $a \in A$,

$$\chi(a) = (\alpha_A(a))(\chi) = (\alpha_{\hat{A}}(\chi))(\alpha_A(a)) = \zeta(\alpha_A(a)) \in \zeta(\alpha_A(A)) = \{0\}. \quad (\text{I.18})$$

Hence, $\chi = 0$ (and so $\zeta = 0$) contrary to our assumption, which completes the proof. $\square$

I.38. Bohr-compactification. The Čech-compactification βX of a Tychonoff space X has the property that every continuous map $f: X \rightarrow K$ into a compact Hausdorff space X extends to βX , and thus factors uniquely through the dense embedding $X \rightarrow \beta X$. It appears to be less known, however, that βX exists even when X is not Tychonoff, in which case the map $X \rightarrow \beta X$ is only continuous but need not be injective or open onto its image. (Every continuous map of X into a compact Hausdorff space still factors uniquely through βX .) In a categorical language, one says that the category HComp of compact Hausdorff spaces and their continuous maps is a reflective subcategory of Top .

Similarly to the relationship between Top and HComp , the full subcategory $\text{Grp}(\text{HComp})$ of compact Hausdorff groups is reflective in $\text{Grp}(\text{Top})$ (topological groups and their continuous homomorphisms). The reflection is called the *Bohr-compactification*, and is denoted by $\rho_G: G \rightarrow bG$. We show its existence for abelian groups, where the construction is rather simple

I.39. Theorem. *For every $A \in \text{Ab}(\text{Top})$ there is $bA \in \text{Ab}(\text{HComp})$ and a continuous homomorphism $\rho_A: A \rightarrow bA$ such that every continuous homomorphism $\varphi: A \rightarrow K$ into $K \in \text{Ab}(\text{HComp})$ factors uniquely through ρ_A :*

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & K \\ & \searrow \rho_A & \nearrow \exists! \tilde{\varphi} \\ & bA & \end{array} \quad (\text{I.19})$$

Moreover, $\widehat{bA} = (\hat{A})_d$, and $\ker \rho_A = \ker \alpha_A$.

Proof. Since bA is supposed to be compact, by Lemma I.36(a), $\widehat{bA}$ must be discrete. Applying the universal property of bA to $K = \mathbb{T}$ and $\varphi = \chi \in \hat{A}$ yields that A and bA have the same continuous characters, that is, $\widehat{bA} = (\hat{A})_d$ (the subscript d stands for the discrete topology). Using this necessary condition as a definition, one set $bA = (\hat{A})_d$ and $(\rho_A(a))(\chi) = \chi(a)$ for every $a \in A$ and $\chi \in (\hat{A})_d$. By Proposition I.5(b), a basic open set in bA has the form $\Phi^\triangleright$, where $\Phi \subseteq \hat{A}$ is finite. Thus, $\rho_A^{-1}(\Phi^\triangleright) = \bigcap_{\chi \in \Phi} \chi^{-1}(\Lambda_1)$ is a neighborhood of 0 in A , and therefore ρ_A is continuous.

If $\varphi: A \rightarrow K$ is a continuous homomorphism, then it induces $\hat{\varphi}: \hat{K} \rightarrow \hat{A}$. Since $\hat{K}$ is discrete (cf. Lemma I.36(a)), $(\hat{\varphi})_d: \hat{K} \rightarrow (\hat{A})_d$ is also continuous, and so $(\widehat{\hat{\varphi}})_d: (\hat{A})_d \rightarrow \hat{K}$ is continuous as well. By Theorem I.37, $\hat{K} \cong K$, and therefore $\tilde{\varphi} = (\widehat{\hat{\varphi}})_d$. This consideration shows the uniqueness of $\tilde{\varphi}$ too, because $\tilde{\varphi}$ is uniquely determined by its dual. Since $\alpha_A(a)$ and $\rho_A(a)$ are both evaluations at a , the last statement follows. $\square$

A subgroup $H \leq A$ such that $H^{\triangleright\triangleleft} = H$ is said to be *dually closed* in A ; if every continuous character of H extends continuously to A , we say that H is *dually embedded* in A . We saw in Theorem I.19(a) that every dense subgroup of a topological group is dually embedded.

I.40. Corollary. *Let $A \in \text{Ab}(\text{Top})$ be such that α_A is injective, and let K be a compact subgroup of A . Then K is dually closed and dually embedded in A .*

Proof. Since α_A is injective, A must be Hausdorff. First, suppose that A is compact. In order to show that K is dually closed, let $x \in A \setminus K$. The quotient A/K is also compact, and the image $\bar{x}$ of x in A/K is non-zero. Thus, by Theorem I.37, there is $\bar{\chi} \in \widehat{A/K}$ such that $\bar{\chi}(\bar{x}) \neq 0$. The character $\bar{\chi}$ induces $\chi \in \hat{A}$ by setting $\chi(a) = \bar{\chi}(\bar{a})$ (where $\bar{a}$ stands for the image of a in A/K), and $K \subseteq \ker \chi$, while $\chi(x) \neq 0$. Therefore, $\chi \in K^\triangleright$, and hence $x \notin K^{\triangleright\triangleleft}$. (Note that by Lemma I.4, $K^\triangleright$ is a subgroup of $\hat{A}$, and so $K^{\triangleright\triangleleft}$ coincides with its annihilator in A .) This shows that $K = K^{\triangleright\triangleleft}$. In order to show that K is dually embedded, consider the discrete group $\hat{A}$. By Lemma I.4(b), $\hat{\pi}_K: \widehat{\hat{A}/K^\triangleright} \rightarrow \hat{A}$ is injective and its image is $K^{\triangleright\triangleright}$. Since $\hat{A} \cong A$ (cf. Theorem I.37), $K^{\triangleright\triangleright} \cong K^{\triangleright\triangleleft} = K$. The quotient $\hat{A}/K^\triangleright$ is discrete, and so $\widehat{\hat{A}/K^\triangleright}$ is compact. Therefore, $\hat{\pi}_K: \widehat{\hat{A}/K^\triangleright} \rightarrow A$ is an embedding onto K , that is, $K \cong \widehat{\hat{A}/K^\triangleright}$. Hence, applying Theorem I.37 to the compact group $\hat{A}/K^\triangleright$ yields $\hat{K} \cong \widehat{\widehat{\hat{A}/K^\triangleright}} \cong \hat{A}/K^\triangleright$. In particular, $\hat{\iota}_A: \hat{A} \rightarrow \hat{K}$ given by restriction to K is surjective, as desired.

In the general case, since ρ_A continuous, $\rho_A(K)$ is a compact subgroup of bA . Therefore, by what we have shown so far, it is dually closed and dually embedded in bA . Thus, for every $x \in bA \setminus \rho_A(K)$, there is $\chi \in \widehat{bA} = (\hat{A})_d$ such that $\chi(\rho_A(K)) = 0$ and $\chi(x) \neq 0$. In particular, this holds for every $x = \rho_A(a)$, where $a \in A \setminus K$. (Since ρ_A is injective, $a \in A \setminus K$ implies $x \in bA \setminus \rho_A(K)$.) Therefore, $\chi \in K^\triangleright$, and so $a \notin K^{\triangleright\triangleleft}$. Hence, $K = K^{\triangleright\triangleleft}$, as desired. For the second statement, observe that $K \cong \rho_A(K)$ (because ρ_A is injective, K is compact, and bA is Hausdorff). Thus, if $\chi \in \hat{K} = \widehat{\rho_A(K)}$, then it admits a continuous extension ψ to bA , because $\rho_A(K)$ is dually embedded in bA . Therefore, $\psi \circ \rho_A \in \hat{A}$ is a continuous extension of χ . Hence, K is dually embedded in A , as desired. $\square$

I.41. Open and compact subgroups. Open subgroups of abelian groups are closely related to compact subgroups of the Pontryagin dual, and vice versa. We turn to establishing the properties of open and compact subgroup of abelian groups. Our presentation of these results was strongly influenced by the work of Banaszczyk, Chasco, Martín-Peinador, and Bruguera (cf. [4] and [7]).

I.42. Proposition. ([4, 2.2]) *Let $A \in \text{Ab}(\text{Top})$, $U \leq A$ be an open subgroup, and consider the exact sequence*

$$0 \longrightarrow U \xrightarrow{\iota_U} A \xrightarrow{\pi_U} A/U \longrightarrow 0. \quad (\text{I.20})$$

- (a) *The map $\hat{\pi}_U: \widehat{A/U} \rightarrow \hat{A}$ is an embedding and $\widehat{A/U} \cong U^\triangleright$.*
- (b) *The subgroup U is dually embedded in A . In other words, $\hat{\iota}_U: \hat{A} \rightarrow \hat{U}$ is surjective.*
- (c) *$\hat{\iota}_U$ is open, and thus a quotient.*

(d) *The induced sequence*

$$0 \longrightarrow \widehat{A/U} \xrightarrow{\hat{\pi}_U} \hat{A} \xrightarrow{\hat{\iota}_U} \hat{U} \longrightarrow 0 \quad (\text{I.21})$$

is exact.

Proof. Note that A/U is discrete, because U is open, and so $\widehat{A/U}$ is compact by Lemma I.36(b).

(a) By Lemma I.4(b), $\hat{\pi}_U$ is injective, and its image is $U^\triangleright$. Thus, $\hat{\pi}_U$ is an embedding, because it is continuous, its domain is compact and its codomain is Hausdorff.

(b) Let $\chi \in \hat{U}$, and let $V \in \mathcal{N}(U)$ be such that $\chi(V) \subseteq \Lambda_1$ (cf. Proposition I.5(a)). Since $\mathbb{T}$ is an injective abelian group, χ admits an extension $\psi: A \rightarrow \mathbb{T}$. Thus, $\psi(V) = \chi(V) \subseteq \Lambda_1$, and $V \in \mathcal{N}(A)$, because U is open in A . Therefore, by Proposition I.5(a), ψ is continuous on A .

(c) In order to make the proof more transparent, we decomposed it into three simple steps.

Step 1: $A = \langle U, b \rangle$ for some $b \in A$. Let l be the order of b in A/U (possibly infinite). Given a compact subset $K \subseteq A$, it can be covered by finitely many translates $k_1b + U, \dots, k_mb + U$ of U , and without loss of generality we may assume that $0 \leq k_i < l$. Since U is an open subgroup, it is also closed, and so $(K - k_ib) \cap U$ is compact. Thus, $C = ((K - k_1b) \cup \dots \cup (K - k_mb)) \cap U$ is a compact subset of U , and $K \subseteq (k_1b + C) \cup \dots \cup (k_mb + C)$. Without loss of generality, we may assume that $lb \in C$ if $l \neq \infty$. Set $W = \{\chi \in \hat{U} \mid \chi(C) \subseteq \Lambda_2\}$, and we show that $W \subseteq \hat{\iota}_U(K^\triangleright)$. To that end, let $\bar{\chi} \in W$. If $\bar{\chi} \in \hat{A}$ is an extension of χ , then

$$\bar{\chi}(K) \subseteq \bar{\chi}(k_1b + C) \cup \dots \cup \bar{\chi}(k_mb + C) \subseteq (k_1\bar{\chi}(b) + \Lambda_2) \cup \dots \cup (k_m\bar{\chi}(b) + \Lambda_2). \quad (\text{I.22})$$

If l is finite, then $lb \in C$, and so $\chi(lb) \in \Lambda_2$. Let $r \in \Lambda_{2l} \subseteq \mathbb{T}$ be the closest point to 0 such that $lr = \chi(lb)$ (if $l = \infty$, then $r = 0$). Set $\bar{\chi}(u + nb) = \chi(u) + nr$. The character $\bar{\chi}$ is continuous on A because $\bar{\chi}|_U = \chi$ is continuous, and U is an open subgroup. Furthermore, one has $k_i\bar{\chi}(b) \in \Lambda_2$ for each i , because $0 \leq k_i < l$ and $r \in \Lambda_{2l}$. Therefore, by (I.22), $\bar{\chi}(K) \subseteq \Lambda_2 + \Lambda_2 \subseteq \Lambda_1$, which means that $\bar{\chi} \in K^\triangleright$. Hence, $W \subseteq \hat{\iota}_U(K^\triangleright)$, as desired.

Step 2: A/U is finitely generated, that is, $A = \langle U, b_1, \dots, b_n \rangle$ for $b_1, \dots, b_n \in A$. Set $U_0 = U$ and $U_k = \langle U_{k-1}, b_k \rangle$. Each U_k is an open subgroup of U_{k+1} (and of A), and thus by step 1, each $\hat{\iota}_{U_k}: \hat{U}_{k+1} \rightarrow \hat{U}_k$ is open. Therefore, $\hat{\iota}_U = \hat{\iota}_{U_{k-1}} \circ \dots \circ \hat{\iota}_{U_0}$ is open.

Step 3: In the general case, let $K \subseteq A$ be compact. Then K can be covered by finitely many translates of U , $K \subseteq (b_1 + U) \cup \dots \cup (b_n + U)$. Put $U' = \langle U, b_1, \dots, b_n \rangle$. Since $K \subseteq U'$, the image $\hat{\iota}_{U'}(K^\triangleright)$ coincides with the polar of K in $\hat{U}'$ with respect to U' , and thus open. By step 2, $\hat{\iota}_U: \hat{U}' \rightarrow \hat{U}$ is open. Therefore, $\hat{\iota}_U(K^\triangleright)$ is open in $\hat{U}$, as desired.

(d) Exactness at $\widehat{A/U}$ and $\hat{U}$ were shown in (b) and (c), respectively. Exactness at $\hat{A}$ also follows from (b), because $\text{Im } \hat{\pi}_U = U^\triangleright = \ker \hat{\iota}_U$. $\square$

I.43. Proposition. Let $A \in \text{Ab}(\text{Top})$, $K \leq A$ be a compact subgroup, and consider the exact sequence

$$0 \longrightarrow K \xrightarrow{\iota_K} A \xrightarrow{\pi_K} A/K \longrightarrow 0. \quad (\text{I.23})$$

(a) *The map $\hat{\pi}_K: \widehat{A/K} \rightarrow \hat{A}$ is an embedding and $\widehat{A/K} \cong K^\triangleright$.*

- (b) The map $\hat{\iota}_K: \hat{A} \rightarrow \hat{K}$ is open onto its image.
(c) The induced sequence

$$0 \longrightarrow \widehat{A/K} \xrightarrow{\hat{\pi}_K} \hat{A} \xrightarrow{\hat{\iota}_K} \hat{K} \quad (\text{I.24})$$

is exact.

- (d) If α_A is injective, then $\hat{\iota}_K: \hat{A} \rightarrow \hat{K}$ is surjective, and so $\hat{\iota}_K$ is a quotient map.

Proof. Note that by Lemma I.36(a), $\hat{K}$ is compact.

(a) A basic neighborhood of 0 in $\widehat{A/K}$ is of the form $L^\triangleright$, where $L \subseteq A/K$ is compact (cf. Proposition I.5(b)). By Lemma A.3(b), $\pi_K^{-1}(L)$ is a compact subset of A , and thus $\pi_K^{-1}(L)^\triangleright$ is a basic neighborhood of 0 in $\hat{A}$. Without loss of generality, we may assume that $0 \in L$, and then by Lemma I.4(c), $\pi_K^{-1}(L)^\triangleright = \hat{\pi}_K(L^\triangleright)$, as desired.

(b) Since $\ker \hat{\iota}_K = \widehat{K^\triangleright}$ is open in $\hat{A}$ and $\hat{K}$ is discrete, the statement is obvious.

(c) Exactness at $\widehat{A/K}$ and at $\hat{A}$ follows from (b), because $\text{Im } \hat{\pi}_K = K^\triangleright = \ker \hat{\iota}_K$.

(d) By Corollary I.40, K is dually embedded in A , which completes the proof. $\square$

I.44. Theorem. ([4, 2.3]) Let $A \in \text{Ab}(\text{Top})$, and let $U \leq A$ be an open subgroup. Then:

- (a) α_U is injective (resp., surjective) if and only if α_A is injective (resp., surjective);
(b) α_U is an isomorphism of topological groups if and only if α_A is so.

Proof. By Proposition I.42, the exact sequence

$$0 \longrightarrow U \xrightarrow{\iota_U} A \xrightarrow{\pi_U} A/U \longrightarrow 0 \quad (\text{I.25})$$

gives rise to an exact sequence

$$0 \longrightarrow \widehat{A/U} \xrightarrow{\hat{\pi}_U} \hat{A} \xrightarrow{\hat{\iota}_U} \hat{U} \longrightarrow 0 \quad (\text{I.26})$$

with $\widehat{A/U}$ compact, $\hat{\pi}_U$ an embedding, and $\hat{\iota}_U$ a quotient map. Thus, the conditions of Proposition I.43 are fulfilled (by Remark I.13(3), $\alpha_{\hat{A}}$ is injective), and so

$$0 \longrightarrow \hat{U} \xrightarrow{\hat{\iota}_U} \hat{A} \xrightarrow{\hat{\pi}_U} \widehat{\widehat{A/U}} \longrightarrow 0 \quad (\text{I.27})$$

is also exact, $\hat{\iota}_U$ is an embedding, and $\hat{\pi}_U$ is a quotient map. Since the evaluation homomorphism α is a natural transformation, we obtain a commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & U & \xrightarrow{\iota_U} & A & \xrightarrow{\pi_U} & A/U \longrightarrow 0 \\ & & \alpha_U \downarrow & & \alpha_A \downarrow & & \alpha_{A/U} \downarrow \\ 0 & \longrightarrow & \hat{U} & \xrightarrow{\hat{\iota}_U} & \hat{A} & \xrightarrow{\hat{\pi}_U} & \widehat{\widehat{A/U}} \longrightarrow 0 \end{array} \quad (\text{I.28})$$

The group A/U is discrete, and thus $\alpha_{A/U}$ is a topological isomorphism by Theorem I.37. Therefore, (a) follows from the well-known Five Lemma for abelian groups. In order to show (b), observe that $\alpha_U = \alpha_A|_U$ and $\alpha_U^{-1} = \alpha_A^{-1}|_U$, and that it suffices to check the continuity of a homomorphism on a neighborhood of 0. Since U is open, the statement follows. $\square$

I.45. Theorem. ([4, 2.6]) Let $A \in \text{Ab}(\text{Haus})$ be such that α_A is injective, and let $K \leq A$ be a compact subgroup. Then:

- (a) $\alpha_{A/K}$ is injective;
- (b) $\alpha_{A/K}$ is surjective if and only if α_A is so;
- (c) $\alpha_{A/K}$ is an isomorphism of topological groups if and only if α_A is so.

I.46. Remark. Although in Proposition I.43 we assumed no separation axioms, here we do, because α_A being injective implies that A is Hausdorff.

Proof. Since α_K is injective, by Proposition I.43, the exact sequence in (I.23) gives rise to an exact sequence

$$0 \longrightarrow \widehat{A/K} \xrightarrow{\hat{\pi}_K} \hat{A} \xrightarrow{\hat{\iota}_K} \hat{K} \longrightarrow 0 \quad (\text{I.29})$$

where $\widehat{A/K}$ is an open subgroup of $\hat{A}$ and $\hat{\iota}_K$ is a quotient. Therefore, by Proposition I.42, the lower row of the commutative diagram below is exact:

$$\begin{array}{ccccccc} 0 & \longrightarrow & K & \xrightarrow{\iota_K} & A & \xrightarrow{\pi_K} & A/K \longrightarrow 0 \\ & & \downarrow \alpha_K & & \downarrow \alpha_A & & \downarrow \alpha_{A/K} \\ 0 & \longrightarrow & \hat{K} & \xrightarrow{\hat{\iota}_K} & \hat{A} & \xrightarrow{\hat{\pi}_K} & \widehat{A/K} \longrightarrow 0 \end{array} \quad (\text{I.30})$$

Since K is compact Hausdorff, α_K is an isomorphism of topological groups (cf. Proposition I.37), and thus (a) and (b) follow from the Five Lemma for abelian groups.

(c) If α_A is an isomorphism of topological groups, then $\alpha_{A/K}$ is a bijection by (a) and (b), and the topologies of A/K and $\widehat{A/K}$ coincide, because $\widehat{A/K} \cong \hat{A}/\hat{K} \cong A/K$.

Conversely, suppose that $\alpha_{A/K}$ is a topological isomorphism. Then α_A is bijective by (a) and (b). We show that α_A is continuous. To that end, let $\mathcal{F}$ be a filter converging to 0 in A . Then $\pi_K(\mathcal{F}) \longrightarrow K$, and so $\hat{\pi}_K(\alpha_A(\mathcal{F})) = \alpha_{A/K}(\pi_K(\mathcal{F})) \longrightarrow \hat{K}$. By Lemma A.3(a), the filter $\alpha_A(\mathcal{F})$ has a cluster point $x \in \hat{K} \subseteq \hat{A}$, because $\hat{K}$ is compact. So, let $\mathcal{G} \supseteq \mathcal{F}$ be a filter such that $\alpha_A(\mathcal{G}) \longrightarrow x$. For $\chi \in \hat{A}$, one has

$$x(\chi) = \lim \alpha_A(\mathcal{G})(\chi) = \lim \chi(\mathcal{G}) = \chi(\lim \mathcal{G}) = 0 \quad (\text{I.31})$$

because χ is continuous and $\mathcal{G} \longrightarrow 0$, and thus $x = 0$. Therefore, 0 is the unique cluster point of $\alpha_A(\mathcal{F})$ for every filter $\mathcal{F} \longrightarrow 0$. Hence, $\alpha_A(\mathcal{F}) \longrightarrow 0$ for every $\mathcal{F} \longrightarrow 0$, and α_A is continuous.

Since the situation is completely symmetric (one could invert α_K , α_A , and $\alpha_{A/K}$), the proof of the continuity of α_A^{-1} is similar. $\square$

I.47. Locally compact groups. We conclude this chapter with making a step toward the classical Pontryagin duality for locally compact Hausdorff abelian (LCA) groups. We denote by LC the category of locally compact Hausdorff spaces, and thus $\text{Ab}(\text{LC})$ is the category of LCA groups.

I.48. Proposition. Let $A \in \text{Ab}(\text{LC})$. Then:

- (a) $\hat{A} \in \text{Ab}(\text{LC})$;
- (b) α_A is continuous;
- (c) α_A is injective.

Proof. (a) Clearly, $\hat{A}$ is Hausdorff. Let $U \in \mathcal{N}(A)$ such that $\bar{U}$ is compact. By Proposition I.7(a), $U^\triangleright$ is compact in $\hat{A}$, and by Proposition I.5(a), $\bar{U}^\triangleright$ is a basic neighborhood of 0 in $\hat{A}$. Since $\bar{U}^\triangleright \subseteq U^\triangleright$, this completes the proof.

(b) Let $V \in \mathcal{N}(A)$ be such that $\bar{V}$ is compact. By Theorem I.9, α_A is k -continuous, and so $\alpha_A|_{\bar{V}}$ is continuous. In particular, α_A is continuous at 0, and therefore it is continuous.

(c) Let $a \in A$ be a non-zero element.

First, suppose that A is generated by $V \in \mathcal{N}(A)$ such that $\bar{V}$ is compact. By replacing V with $V \cup (-V) \cup \{a, -a\}$ if necessary, we may assume that $a \in V$ and $-V = V$ from the outset. By [33, Lemma 2.42], there is a subgroup $H \leq A$ such that $H \cap V = \{0\}$, $H \cong \mathbb{Z}^n$ for some $n \in \mathbb{N}$, and A/H is compact. The image $a + H$ of a in A/H is non-zero, because $a \in V$ and $V \cap H = \{0\}$. Thus, by Theorem I.37(b), $\alpha_{A/H}(a + H) \neq 0$, and so there is $\bar{\chi} \in \widehat{A/H}$ such that $\bar{\chi}(a + H) \neq 0$. Therefore, one has $\alpha_A(a)(\chi) = \chi(a) = \bar{\chi}(a + H) \neq 0$, where $\chi = \hat{\pi}_H(\bar{\chi}) \in \hat{A}$. Hence, $\alpha_A(a) \neq 0$.

In the general case, pick $V \in \mathcal{N}(A)$ such that $-V = V$, $a \in V$, and $\bar{V}$ is compact, and put $U = \langle V \rangle$, the subgroup generated by V . Then $U + V \subseteq U$, and thus $\bar{U} \subseteq \text{Int } U$ (by Proposition A.1(a)). Therefore, U is an open subgroup of A . By what we proved so far, α_U is injective. Hence, by Theorem I.44(a), α_A is injective. $\square$

I.49. Remark. The proof of (c) falls short of being self-contained because it is based on a Lemma from [33]. I hope to find a simple proof of the relevant parts of the Lemma. Suggestions???

I.50. Lemma. Let $A \in \text{Ab}(\text{LC})$, and suppose that there is a countable family $\{U_n\} \subseteq \mathcal{N}(A)$ of neighborhoods of 0 in A such that $\bigcap U_n = \{0\}$. Then A is metrizable.

Proof. By replacing each U_n with $U'_n \in \mathcal{N}(A)$ such that $\bar{U}'_n \subseteq U_n$ and $\bar{U}'_n$ is compact if necessary, we may assume that $\bigcap \bar{U}_n = \{0\}$, and $\bar{U}_1$ is compact (such U'_n exists, because A is regular). Furthermore, without loss of generality, we may assume that $U_{n+1} \subseteq U_n$. By Theorem I.29, it suffices to show that $\{U_n\}$ is a base at 0 for A . To that end, let $V \in \mathcal{N}(A)$. Then $\bigcap (\bar{U}_n \setminus \text{Int } V) = \emptyset$, and $\{\bar{U}_n \setminus \text{Int } V\}$ is a decreasing family of closed subsets of the compact space $\bar{U}_1$. Therefore, there is n_0 such that $\bar{U}_{n_0} \setminus \text{Int } V = \emptyset$. Hence, $\bar{U}_{n_0} \subseteq \text{Int } V$, as desired. $\square$

I.51. Lemma. Let $A \in \text{Ab}(\text{LC})$, and let $V \in \mathcal{N}(A)$ be such that $\bar{V}$ is compact. Then V contains a compact subgroup K of A such that A/K is metrizable.

Proof. We construct a family $\{V_n\}$ of neighborhoods of 0 in A . Put $V_1 = (-V) \cap V$, and for $n \in \mathbb{N}$, using continuity of addition in A , pick $V_{n+1} \in \mathcal{N}(A)$ such that $V_{n+1} + V_{n+1} \subseteq V_n$ and $-V_{n+1} = V_{n+1}$. Put $K = \bigcap_{n=1}^{\infty} V_n$; K is a subgroup of A , because $K - K \subseteq V_{n+1} - V_{n+1} \subseteq V_n$ implies $K - K \subseteq K$. One has $K + V_{n+1} \subseteq V_{n+1} + V_{n+1} \subseteq V_n$, and thus $\bar{K} \subseteq \text{Int } V_n$ (cf. Proposition A.1(a)). Therefore, $\bar{K} \subseteq K \subseteq \bar{V}$, and K is compact. The inclusion $K + V_{n+1} \subseteq V_n$ also yields $K = \bigcap_{n=1}^{\infty} (K + V_n)$, which means that $\bigcap_{n=1}^{\infty} \pi_K(V_n) = K$ in A/K . Hence, $\{\pi_K(V_n)\}$ satisfies the conditions of Lemma I.50, and A/K is metrizable, as desired. $\square$

I.52. Theorem. The following statements are equivalent:

- (i) For every second countable $A \in \text{Ab}(\text{LC})$, α_A is an isomorphism of topological groups.
(ii) For every $A \in \text{Ab}(\text{LC})$, α_A is an isomorphism of topological groups.

Proof. (i) $\Rightarrow$ (ii): Let $A \in \text{Ab}(\text{LC})$. First, suppose that A is generated by $V \in \mathcal{N}(A)$ such that $\bar{V}$ is compact. By replacing V with $-V \cup V$ if necessary, we may assume that $-V = V$ from outset. Then $A = \bigcup_{n=1}^{\infty} \underbrace{\bar{V} + \cdots + \bar{V}}_{n \text{ times}}$, so A is σ -compact, and in particular, A is Lindelöf. Let

$K \subseteq V$ be a compact subgroup such that A/K is metrizable (cf. Lemma I.51). The quotient A/K is second countable, because it is a continuous image of the Lindelöf space A . (The properties Lindelöf, separable, and second countable are equivalent for metrizable spaces.) Thus, $\alpha_{A/K}$ is an isomorphism of topological groups by our assumption. By Proposition I.48(c), α_A is injective, and therefore, by Theorem I.45(c), α_A is an isomorphism of topological groups.

In the general case, pick $V \in \mathcal{N}(A)$ such that $\bar{V}$ is compact, and put $U = \langle V \rangle$, the subgroup generated by V . Then $U + V \subseteq U$, and thus $\bar{U} \subseteq \text{Int } U$ (by Proposition A.1(a)). Therefore, U is an open subgroup of A . By what we proved so far, α_U is an isomorphism of topological groups, because U is generated by a compact neighborhood of 0. Hence, by Theorem I.44(b), α_A is an isomorphism of topological groups. $\square$

Appendix

A. Separation properties of topological groups

For $G \in \text{Grp}(\text{Top})$, one puts $\mathcal{N}(G)$ for the collection of neighborhoods of e in G .

A.1. Proposition. *Let $G \in \text{Grp}(\text{Top})$.*

- (a) *If $VW \subseteq U$ for $U, V, W \in \mathcal{N}(G)$, then $\bar{V}, \bar{W} \subseteq \text{Int}U$.*
- (b) *G is regular.*
- (c) *The following statements are equivalent:*
 - (i) *G is T_3 (i.e., T_1 and regular);*
 - (ii) *G is Hausdorff;*
 - (iii) *G is T_1 ;*
 - (iv) *G is T_0 .*

Proof. (a) Set $W_1 = (\text{Int } W)^{-1}$, and let $x \in \bar{V}$. Then $W_1 \in \mathcal{N}(G)$ (because the group inversion is continuous), and $V \cap xW_1 \neq \emptyset$. Thus, there is $v \in W$ and $w_1 \in W_1$ such that $xw_1 = v$, or $x = vw_1^{-1}$. Therefore, $x \in VW_1^{-1} = V(\text{Int } W)$. Since $V(\text{Int } W)$ is open, it is contained in $\text{Int}(VW) \subseteq \text{Int } U$. Hence, $x \in \text{Int } U$, as desired.

(b) By continuity of the multiplication group $m: G \times G \rightarrow G$, for every $U \in \mathcal{N}(G)$ there is $V \in \mathcal{N}(G)$ such that $VV \subseteq U$, and thus $\bar{V} \subseteq \text{Int } U$ by (a).

(c) Implications (i) $\Rightarrow$ (ii) $\Rightarrow$ (iii) $\Rightarrow$ (iv) are obvious. In order to complete the proof, observe that by (b), G is regular, and every T_0 regular topological space X is T_1 . (Indeed, if $x, y \in X$ are distinct points, then one of them, say x , has a neighborhood U_0 such that $y \notin U_0$. Then $F = X \setminus \text{Int } U_0$ is a closed subset containing y that does not contain x . Since X is regular, there are disjoint open subsets U and V of X such that $F \subseteq U$ and $x \in V$. This completes the proof.) $\square$

A.2. Proposition. *Let $G \in \text{Grp}(\text{Top})$, and set $N_G = \bigcap \mathcal{N}(G)$. Then:*

- (a) *N_G is a compact normal subgroup of G ;*
- (b) *G/N_G is Hausdorff;*
- (c) *for every continuous homomorphism $\varphi: G \rightarrow H$ into a Hausdorff group H , $N_G \subseteq \ker \varphi$. In particular, φ factors uniquely through $G \rightarrow G/N_G$.*

Proof. (a) Let $U \in \mathcal{N}(G)$ and $g \in G$. By continuity of the group operations, there is $V \in \mathcal{N}(G)$ such that $VV^{-1} \subseteq U$ and $g^{-1}Vg \subseteq U$. Thus, $N_G N_G^{-1} \subseteq VV^{-1} \subseteq U$ and $g^{-1}N_G g \subseteq g^{-1}Vg \subseteq U$. This is true for every $U \in \mathcal{N}(G)$, and therefore $N_G N_G^{-1} \subseteq N_G$ and $g^{-1}N_G g \subseteq N_G$. Hence, N_G is a normal subgroup. Since N_G is contained in every neighborhood of e , its compactness is clear.

(b) For every $U \in \mathcal{N}(G)$, there is $V \in \mathcal{N}(G)$ such that $VV \subseteq U$, and thus $VN_G \subseteq VV \subseteq U$. Thus, $N_G \subseteq \bigcap \{VN_G \mid V \in \mathcal{N}(G)\} \subseteq \bigcap \mathcal{N}(G) = N_G$. Therefore, G/N_G is T_0 , and hence it is Hausdorff, by Proposition A.1.

(c) If H is Hausdorff, then $\{e_H\} = \bigcap \mathcal{N}(H)$, and so

$$\ker \varphi = \bigcap \{\varphi^{-1}(W) \mid W \in \mathcal{N}(H)\} \supseteq \bigcap \mathcal{N}(G) = N_G, \quad (1)$$

as desired. $\square$

The group G/N_G is the *maximal Hausdorff quotient* of G . In categorical language, this means that G/N_G is the reflection of $G \in \text{Grp}(\text{Top})$ in $\text{Grp}(\text{Haus})$.

A.3. Lemma. *Let $G \in \text{Grp}(\text{Top})$, $K \leq G$ be a compact subgroup, and put $\pi_K: G \rightarrow G/K$ for the canonical projection.*

(a) *If $\mathcal{F}$ is a filter in G such that K is a cluster point of $\pi_K(\mathcal{F})$ in G/K , then $\mathcal{F}$ has a cluster point in K .*

(b) *If $L \subseteq G/K$ is compact, then so is $\pi_K^{-1}(L)$.*

Proof. (a) Assume that $\mathcal{F}$ has no cluster point in K . Then each point $x \in K$ has a neighborhood U_x such that U_x does not mesh $\mathcal{F}$, that is, there is $F_x \in \mathcal{F}$ such that $U_x \cap F_x = \emptyset$. The collection $\{U_x\}_{x \in K}$ is an open cover of K , and thus it has a finite subcover $U_{x_1}, \dots, U_{x_l}$, because K is compact. Set $U = U_{x_1} \cup \dots \cup U_{x_l}$ and $F = F_{x_1} \cap \dots \cap F_{x_l}$. Since $\mathcal{F}$ is a filter, $F \in \mathcal{F}$, and one has $U \cap F = \emptyset$ and $K \subseteq U$. Using the group multiplication $m: G \times G \rightarrow G$, the last inclusion can be expressed as $\{e\} \times K \subseteq m^{-1}(U)$. Since m is continuous, $m^{-1}(U)$ is open, and thus, by a tube-lemma type argument, there is $V \in \mathcal{N}(G)$ such that $V \times K \subseteq m^{-1}(U)$. In other words, $VK \subseteq U$. Therefore, $VK \cap F = \emptyset$, and so $VK \cap FK = \emptyset$. Hence, $\pi_K(V) \cap \pi_K(F) = \emptyset$. Since $\pi_K(V)$ is a neighborhood of K in G/K , this contradicts the assumption that K is a cluster point of $\pi_K(\mathcal{F})$.

(b) Let $\mathcal{F}$ be a filter meshing $\pi_K^{-1}(L)$ (i.e., $F \cap \pi_K^{-1}(L) \neq \emptyset$ for every $F \in \mathcal{F}$). Then $\pi_K(\mathcal{F})$ is a filter that meshes L , and thus has a cluster point xK , because L is compact. Equivalently, K is a cluster point of $\pi_K(x^{-1}\mathcal{F})$, and therefore $x^{-1}\mathcal{F}$ has a cluster point in K (by (a)). Hence, $\mathcal{F}$ has a cluster point in xK , which shows that $\pi_K^{-1}(L)$ is compact. $\square$

A combination of Proposition A.2(a) and Lemma A.3 yields:

A.4. Corollary. *Compact subsets of G/N_G are precisely the images of compact subsets of G .* $\square$

B. Exponentiability and the compact-open topology

Let $\mathfrak{T}$ be a full subcategory of Top , the category of topological spaces and their continuous maps. Suppose that $\mathfrak{T}$ has finite products; they may differ from the products in Top , but each product in $\mathfrak{T}$ must have the same underlying set as in Top . Each function $f: X \times Y \rightarrow Z$ gives rise to a map $\bar{f}: Y \rightarrow Z^X$ defined by $\bar{f}(y) = f_y$, where $f_y \in Z^X$ is given by $f_y(x) = f(x, y)$. One would like to equip the function set $\mathfrak{T}(X, Z) \subseteq Z^X$ with a suitable topology such that $\bar{f}$ is continuous whenever

f is so. If we add the natural requirement of functoriality of the topology, we arrive at the concept of exponentiability. A space $X \in \mathfrak{T}$ is *exponentiable* (in $\mathfrak{T}$) if the functor $X \times - : \mathfrak{T} \longrightarrow \mathfrak{T}$ has a right adjoint. Since we assume $\mathfrak{T}$ to have finite products, it has a terminal object which must then be the one-point space $\{*\}$, because $\mathfrak{T}$ is a full subcategory. Thus, one expects $\mathfrak{T}(X, Z)$ to be the underlying set of the hom-object. In other words, X is exponentiable if there is a way of topologizing the set $\mathfrak{T}(X, Z)$ for every $Z \in \mathfrak{T}$ such that the resulting space is in $\mathfrak{T}$, and the map

$$\begin{aligned} \mathfrak{T}(X \times Y, Z) &\longrightarrow \mathfrak{T}(Y, \mathfrak{T}(X, Z)) \\ f &\longmapsto \bar{f} \end{aligned} \tag{2}$$

is a bijection that is natural in Y and Z .

The question of describing exponentiable Hausdorff spaces was raised by Hurewicz in a personal communication with Fox, who was the first to give a partial answer to the question (cf. [14]). He proved that for every regular locally compact space X and every $Y \in \text{Haus}$ the set $\text{Haus}(X, Y)$ can be topologized in a way that f is continuous if and only if $\bar{f}$ is so. Fox also showed that for every separable metric space X , it is possible to topologize $\text{Haus}(X, \mathbb{R})$ in a way that (2) holds for every Y if and only if X is locally compact. Although Fox did not prove that the condition of local compactness is sufficient, and he showed necessity only for separable metric spaces X , he was actually very close to a complete solution.

For Hausdorff spaces X and Y , we denote by $\mathcal{C}(X, Y)$ the space $\text{Haus}(X, Y)$ equipped with the *compact-open topology*: Its subbase is the family $\{[K, U] \mid K \subseteq X \text{ compact}, U \subseteq Y \text{ open}\}$, where $[K, U] = \{f \in \text{Haus}(X, Y) \mid f(K) \subseteq U\}$. The compact-open topology was also invented by Fox, and following his work, Arens studied the separation properties of $\mathcal{C}(X, Y)$ (cf. [1]). Arens was not far from proving that local compactness is necessary for exponentiability in Haus (cf. [1, Theorem 3]). Jackson proved that if X is locally compact, then $\mathcal{C}(X, -)$ is the right adjoint of $X \times -$, moreover, the bijection

$$\mathcal{C}(X \times Y, Z) \longleftrightarrow \mathcal{C}(Y, \mathcal{C}(X, Z)) \tag{3}$$

is actually a homeomorphism for every Hausdorff space Y and Z (cf. [19]). It appears that Fox already recognized that (3) is a bijection, and Jackson's main achievement is proving that it is a homeomorphism.

B.1. Remark. Some authors, including Isbell, erroneously credit Brown for proving (3) (cf. [18]). The fact is that Brown himself refers both to Fox and Jackson in his paper in question, and he lays no claim to the “classical” results on special cases of the exponential law (as a personal communication with him reveals). At the same time, Brown was the first to show that the category of Hausdorff k -spaces is cartesian closed, but unfortunately this seems to be somewhat forgotten (cf. [5, 3.3]).

Whitehead proved that if X is a locally compact Hausdorff space, then $1_X \times g$ is a quotient map in Haus for every quotient map g in Haus (cf. [35]). Michael proved that the converse is also true: For a Hausdorff space X , the map $1_X \times g$ is a quotient map in Haus for every quotient map g in Haus if and only if X is locally compact (cf. [25, 2.1]). In other words, the functor $X \times -$ preserves quotients if and only if X is locally compact. If X is exponentiable, then $X \times -$ must preserve

every colimit in Haus ([24, V.5.1]), and in particular it has to preserve coequalizers in Haus (which is equivalent to preservation of quotients; for details see [22, 1.1]). Thus, by Michael's result the local compactness of X follows. Therefore, exponentiable spaces in Haus can be characterized as follows.

B.2. Theorem. *A space $X \in \text{Haus}$ is exponentiable in Haus if and only if X is locally compact.*

A proof of sufficiency of local compactness in Theorem B.2 is available in standard topology textbooks (cf. [13, 3.2]).

B.3. Corollary. *Let $X, Y \in \text{Haus}$. The evaluation map*

$$e : X \longrightarrow \mathcal{C}(\mathcal{C}(X, Y), Y) \quad (4)$$

$$x \longmapsto \hat{x} \quad [\hat{x}(f) = f(x)] \quad (5)$$

is k -continuous, that is, $e|_K$ is continuous for every compact subset $K \subseteq X$.

Proof. Let $K \subseteq X$ be a compact subset. The map

$$E : K \times \mathcal{C}(X, Y) \longrightarrow Y \quad (6)$$

$$(x, f) \longmapsto f(x) \quad (7)$$

is continuous, because $E(K \times [K, W]) \subseteq W$ for every open subset W of Y . Let $[\Phi, U]$ be a subbasic open subset of $\mathcal{C}(\mathcal{C}(X, Y), Y)$, where $\Phi \subseteq \mathcal{C}(X, Y)$ is compact and $U \subseteq Y$ is open. By Theorem B.2, Φ is exponentiable (because it is compact), and thus $E|_{K \times \Phi} : K \times \Phi \rightarrow Y$ gives rise to a continuous map $e_{K, \Phi} : K \rightarrow \mathcal{C}(\Phi, Y)$ that is defined by $(e_{K, \Phi}(k))(f) = f(k)$. Therefore, $(e|_K)^{-1}([\Phi, U]) = e_{K, \Phi}^{-1}([\Phi, U])$ is open in K , as desired. $\square$

Exponentiable spaces in Top were characterized by Day and Kelly as so-called *core-compact* spaces (cf. [11]), but this result is beyond the scope of this summary. For a detailed review of exponentiability, we refer the reader to [22, Chapter 1].

C. Hausdorff k -spaces

A map $f : X \rightarrow Y$ between Hausdorff spaces is said to be *k -continuous* if the restriction $f|_K$ is continuous for every compact subset K of X . Theorem I.9 suggests that it might be helpful to consider spaces whose k -continuous maps are continuous.

C.1. Proposition. ([13, 3.3.18-21]) *Let $X \in \text{Haus}$. The following statements are equivalent:*

- (i) *if $O \cap K$ is open in K for every $K \in \mathcal{K}(X)$, then O is open in X ;*
- (ii) *if $F \cap K$ is closed for every $K \in \mathcal{K}(X)$, then F is closed;*
- (iii) *every k -continuous map $f : X \rightarrow Y$ into a Hausdorff space Y is continuous.*

Proof. The equivalence of (i) and (ii) is obvious, and clearly each of them implies (iii).

(iii) $\Rightarrow$ (i): Let kX be the underlying set of X equipped with the topology

$$\{O \subseteq X \mid O \cap K \text{ is open in } K \text{ for every } K \in \mathcal{K}(X)\}. \quad (8)$$

We call kX the *k-ification* of X . Its topology is finer than the topology of X , and so kX is Hausdorff. Furthermore, the topologies of X and kX coincide on compact subsets, and thus the identity map $X \rightarrow kX$ is *k*-continuous. Therefore, by (iii), it is continuous, and hence $X \cong kX$, as desired. $\square$

A Hausdorff space X is a *k-space* if it satisfies the equivalent conditions of Proposition C.1. The category of Hausdorff *k*-spaces and their continuous maps is denoted by $k\text{Haus}$. It is possible to define *k*-spaces without assuming any separation axioms (cf. [34] and [22, 1.1]), but it is unnecessary for our purposes.

C.2. Theorem.

- (a) *The category $k\text{Haus}$ is a coreflective subcategory of Haus with coreflector k , the *k-ification*.*
- (b) *The product of $X, Y \in k\text{Haus}$ in $k\text{Haus}$ is the *k-ification* $k(X \times Y)$ of their product in Haus .*
- (c) ([5, 3.3]) *The category $k\text{Haus}$ is cartesian closed, and the internal hom is given by the *k-ification* $k\mathcal{C}(X, Y)$ of the compact-open topology.*

Proof. (a) First, note that by Proposition C.1, kX is a Hausdorff *k*-space for every $X \in \text{Haus}$. Let $g: X \rightarrow Y$ be a continuous map between Hausdorff spaces X and Y , and let $O \subseteq kY$ be open. Since g is continuous, $g(K) \in \mathcal{K}(Y)$ for every $K \in \mathcal{K}(X)$. Thus, $g(K) \cap O$ is open in $g(K)$, and hence $K \cap g^{-1}(O) = g^{-1}(g(K) \cap O)$ is open in K . Therefore, $kg: kX \rightarrow kY$ is continuous. This shows that $k: \text{Haus} \rightarrow k\text{Haus}$ is a functor (it is clear that it preserves id_X and composition). In order to show that k is a coreflector, observe that if $X \in k\text{Haus}$ (i.e., $X = kX$), then $kf: X \rightarrow kY$ is the unique continuous map such that $f = r_Y \circ kf$, where $r_Y: kY \rightarrow Y$ is the identity.

(b) follows from (a) and [24, VI.3, V.5.1].

It is not hard to see that (c) follows from Corollary B.3, but for a detailed proof, we refer the reader to [24, VII.8.3]. $\square$

Bibliography

- [1] Richard F. Arens. A topology for spaces of transformations. *Ann. of Math. (2)*, 47:480–495, 1946.
- [2] L. Aussenhofer. Contributions to the duality theory of abelian topological groups and to the theory of nuclear groups. *Dissertationes Math. (Rozprawy Mat.)*, 384:113, 1999.
- [3] Wojciech Banaszczyk. *Additive subgroups of topological vector spaces*, volume 1466 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1991.
- [4] Wojciech Banaszczyk, María Jesús Chasco, and Elena Martín Peinador. Open subgroups and Pontryagin duality. *Math. Z.*, 215(2):195–204, 1994.
- [5] R. Brown. Function spaces and product topologies. *Quart. J. Math. Oxford Ser. (2)*, 15:238–250, 1964.
- [6] M. Montserrat Bruguera. Some properties of locally quasi-convex groups. In *Proceedings of the First Ibero-American Conference on Topology and its Applications (Benicassim, 1995)*, volume 77, pages 87–94, 1997.
- [7] M. Montserrat Bruguera and Elena Martín-Peinador. Open subgroups, compact subgroups and Binz-Butzmann reflexivity. *Topology Appl.*, 72(2):101–111, 1996.
- [8] Montserrat Bruguera and Elena Martín-Peinador. Banach-Dieudonné theorem revisited. *J. Aust. Math. Soc.*, 75(1):69–83, 2003.
- [9] H.-P. Butzmann. Duality theory for convergence groups. In *Proceedings of the International School of Mathematics “G. Stampacchia” (Erice, 1998)*, volume 111, pages 95–104, 2001.
- [10] M. J. Chasco. Pontryagin duality for metrizable groups. *Arch. Math. (Basel)*, 70(1):22–28, 1998.
- [11] B. J. Day and G. M. Kelly. On topologically quotient maps preserved by pullbacks or products. *Proc. Cambridge Philos. Soc.*, 67:553–558, 1970.
- [12] Dikran N. Dikranjan, Ivan R. Prodanov, and Luchezar N. Stoyanov. *Topological groups*, volume 130 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker Inc., New York, 1990. Characters, dualities and minimal group topologies.

- [13] Ryszard Engelking. *General topology*, volume 6 of *Sigma Series in Pure Mathematics*. Heldermann Verlag, Berlin, second edition, 1989. Translated from the Polish by the author.
- [14] Ralph H. Fox. On topologies for function spaces. *Bull. Amer. Math. Soc.*, 51:429–432, 1945.
- [15] Salvador Hernández. Pontryagin duality for topological abelian groups. *Math. Z.*, 238(3):493–503, 2001.
- [16] Edwin Hewitt and Kenneth A. Ross. *Abstract harmonic analysis*. Academic Press Inc., Publishers, New York, 1963.
- [17] J. R. Isbell. *Uniform spaces*. American Mathematical Society, Providence, R.I., 1964.
- [18] John Isbell. General function spaces, products and continuous lattices. *Math. Proc. Cambridge Philos. Soc.*, 100(2):193–205, 1986.
- [19] James R. Jackson. Spaces of mappings on topological products with applications to homotopy theory. *Proc. Amer. Math. Soc.*, 3:327–333, 1952.
- [20] John L. Kelley. *General topology*. D. Van Nostrand Company, Inc., Toronto-New York-London, 1955.
- [21] W. F. Lamartin. On the foundations of k -group theory. *Dissertationes Math. (Rozprawy Mat.)*, 146:32, 1977.
- [22] Gábor Lukács. *c-Compactnes and Generalized Dualities of Topological Groups*. PhD thesis, York University, 2003. <http://at.yorku.ca/p/a/a/o/41.htm>.
- [23] Gábor Lukács. On homomorphism spaces of metrizable groups. *J. Pure Appl. Algebra*, 182(2-3):263–267, 2003.
- [24] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.
- [25] Ernest Michael. Local compactness and Cartesian products of quotient maps and k -spaces. *Ann. Inst. Fourier (Grenoble)*, 18(fasc. 2):281–286 vii (1969), 1968.
- [26] Sidney A. Morris. *Pontryagin duality and the structure of locally compact abelian groups*. Cambridge University Press, Cambridge, 1977. London Mathematical Society Lecture Note Series, No. 29.
- [27] J. Neumann and E. P. Wigner. Minimally almost periodic groups. *Ann. of Math. (2)*, 41:746–750, 1940.
- [28] N. Noble. k -groups and duality. *Trans. Amer. Math. Soc.*, 151:551–561, 1970.
- [29] E. G. Ostling and A. Wilansky. Locally convex topologies and the convex compactness property. *Proc. Cambridge Philos. Soc.*, 75:45–50, 1974.

- [30] L. S. Pontryagin. *Selected works. Vol. 2.* Gordon & Breach Science Publishers, New York, third edition, 1986. Topological groups, Edited and with a preface by R. V. Gamkrelidze, Translated from the Russian and with a preface by Arlen Brown, With additional material translated by P. S. V. Naidu.
- [31] David W. Roeder. Functorial characterizations of Pontryagin duality. *Trans. Amer. Math. Soc.*, 154:151–175, 1971.
- [32] David W. Roeder. Category theory applied to Pontryagin duality. *Pacific J. Math.*, 52:519–527, 1974.
- [33] Walter Rudin. *Fourier analysis on groups.* Wiley Classics Library. John Wiley & Sons Inc., New York, 1990. Reprint of the 1962 original, A Wiley-Interscience Publication.
- [34] Rainer M. Vogt. Convenient categories of topological spaces for homotopy theory. *Arch. Math. (Basel)*, 22:545–555, 1971.
- [35] J. H. C. Whitehead. Note on a theorem due to Borsuk. *Bull. Amer. Math. Soc.*, 54:1125–1132, 1948.